

Cytokine expression and secretion by skeletal muscle cells: regulatory mechanisms and exercise effects

Jonathan M. Peake¹, Paul Della Gatta², Katsuhiko Suzuki³, David C. Nieman⁴

¹ School of Biomedical Sciences and Institute of Health and Biomedical Sciences, Queensland University of Technology, Brisbane, Australia

² Centre for Physical Activity and Nutrition, School of Exercise and Nutrition Sciences, Deakin University, Melbourne, Australia

³ Faculty of Sport Sciences, Waseda University, Cooperative Major in Advanced Health Science, Tokyo University of Agriculture and Technology/Waseda University, Tokorozawa, Saitama, Japan

⁴ Human Performance Laboratory, Appalachian State University, North Carolina Research Campus, Kannapolis, North Carolina, USA

ABSTRACT

Cytokines are important mediators of various aspects of health and disease, including appetite, glucose and lipid metabolism, insulin sensitivity, skeletal muscle hypertrophy and atrophy. Over the past decade or so, considerable attention has focused on the potential for regular exercise to counteract a range of disease states by modulating cytokine production. Exercise stimulates moderate to large increases in the circulating concentrations of interleukin (IL)-6, IL-8, IL-10, IL-1 receptor antagonist, granulocyte-colony stimulating factor, and smaller increases in tumor necrosis factor- α , monocyte chemoattractant protein-1, IL-1 β , brain-derived neurotrophic factor, IL-12p35/p40 and IL-15. Although many of these cytokines are also expressed in skeletal muscle, not all are released from skeletal muscle into the circulation during exercise. Conversely, some cytokines that are present in the circulation are not expressed in skeletal muscle after exercise. The reasons for these discrepant cytokine responses to exercise are unclear. In this review, we address these uncertainties by summarizing the capacity of skeletal muscle cells to produce cytokines, analyzing other potential cellular sources of circulating cytokines during exercise, and discussing the soluble factors and intracellular signaling pathways that regulate cytokine synthesis (e.g., RNA-binding proteins, microRNAs, suppressor of cytokine signaling proteins, soluble receptors).

adhesion molecules, which require direct cell-to-cell contact, and hormones, which are produced by specialized endocrine organs and circulate throughout the body to exert their actions. Most cytokines are inducible mediators, are transported through the systemic circulation and are synthesized rapidly by multiple cell types in response to various stimuli. Individual cell types can express and secrete several cytokines simultaneously in response to a single stimulus (200). Cytokines are pleiotropic because they influence several cell types, and elicit different effects, depending on the type of target cells. They exert their pleiotropic actions in two phases. First, they bind to specific receptors expressed on cells with different origins and/or functions. Second, they mediate signal transduction through various intracellular messengers and transcription factors. The biological effects of cytokines depend on the presence and concentrations of other cytokines with synergistic, additive or counter-regulatory actions (200). Cytokines can act in an autocrine, paracrine or endocrine fashion to induce or suppress their own synthesis and regulate the production of other cytokines and their receptors. They possess an important characteristic of self-limiting synthesis through various auto-regulatory mechanisms (e.g., RNA instability) and negative feedback pathways. These pathways include synthesis of eicosanoids and corticosteroid hormones, expression of soluble receptors, and induction of intracellular transcription factors that block signal transduction (200).

INTRODUCTION

Cytokines comprise a large family of polypeptides or proteins. This family includes interleukins, interferons, growth- and colony-stimulating factors, chemokines, members of the tumor necrosis factor group and transforming growth factors. Cytokines play an integrative and regulatory role as universal intercellular messengers. Once secreted, they can mediate intercellular communication locally or systemically. Alternatively, they can mediate intercellular contact even when bound to cell membranes (200). These characteristics distinguish cytokines from other intercellular messengers such as

The main function of cytokines is to regulate immune function. However, their wide-ranging effects on cell proliferation, differentiation, migration, survival and apoptosis allow them to play a role in homeostatic control of various tissues, organs and systems. For example, together with hormones and neuropeptides, cytokines mediate interactions between the nervous, endocrine and immune systems. Some cytokines also control body temperature, fatigue, appetite and metabolism. A link between cytokines and skeletal muscle was first established almost 50 years ago when researchers identified that an endogenous pyrogen was present in skeletal muscle (203). Cannon and Kluger (32) subsequently made another important discovery that endurance exercise induces the systemic release of a pyrogenic compound. These findings have since stimulated considerable interest in the biological significance and regulation of cytokine production in muscle during exercise and as a result of sepsis, aging, cancer cachexia and chronic inflammatory diseases.

Contact details:

Jonathan Peake, Institute of Health and Biomedical Innovation, Queensland University of Technology, Brisbane, QLD 4059, Australia
Ph: +61 7 3138 6140, Email: jonathan.peake@qut.edu.au

Initially, exercise research focused on the role of cytokines in mediating inflammatory responses to exercise-induced muscle damage. Evidence has accumulated over the past decade that cytokines play a much broader role during exercise. We now know that cytokines act in a hormone-like manner during exercise, mediating metabolism in working skeletal muscle, the liver and adipose tissue, angiogenesis and neurobiology (166). Following exercise, there is an increase in the circulating concentrations of assorted cytokines. Gene expression for some of these cytokines also increases within skeletal muscle. However, there appears to be a dissociation between local gene expression in skeletal muscle and the systemic concentration of other cytokines. For example, after exercise, tumor necrosis factor (TNF)- α and interleukin (IL)-1 β expression in skeletal muscle increases, but the circulating concentration of these cytokines does not change (or only increases slightly). Conversely, the circulating concentrations of IL-1 receptor antagonist (IL-1ra) and IL-10 increase markedly, but these cytokines are not expressed in skeletal muscle after exercise.

Several excellent reviews have discussed the evidence that skeletal muscle is a secretory organ (167, 168, 183, 232). Yet relatively few studies have specifically examined whether skeletal muscle cells themselves are the main source of cytokine gene expression in skeletal muscle during exercise. In this review, we examine this notion in more detail by (1) reviewing the findings from studies of cytokine expression and secretion in cultured skeletal muscle cells, and (2) summarizing the results of studies that have used histological staining of cytokine expression in cross-sections of muscle tissue. We also discuss other potential sources of cytokines

both in skeletal muscle, other tissues and the systemic circulation. Most research to date has investigated the systemic factors and intracellular signaling pathways that stimulate cytokine secretion by skeletal muscle cells. By contrast, much less is known about the factors that restrict or inhibit cytokine translation in skeletal muscle cells, and cytokine release into the circulation during exercise. We propose some potential negative regulatory mechanisms that may govern cytokine expression and secretion by skeletal muscle cells. Considering the important role of cytokines as local and systemic mediators of various aspects of health and disease, we contend that continuing research is needed to determine the dominant sources and regulation of cytokine production in the body.

LOCAL AND SYSTEMIC CYTOKINE RESPONSES TO EXERCISE

Numerous studies have investigated changes in the circulating concentrations of cytokines following exercise. Cytokine responses are generally dependent on the combination of mode, intensity, and duration of exercise. In the case of IL-6, prolonged running produces the greatest increase in plasma IL-6 concentration (167, 184). Indeed, circulating IL-6 can increase up to 120 \times following endurance exercise. IL-1ra (up to 90 \times), IL-10 (up to 80 \times), IL-8 (15 \times) and monocyte chemoattractant protein (MCP-1) (up to 3 \times) also consistently increase in the circulation following exercise (39, 64, 68, 102, 151-154, 164, 165, 208, 212, 219, 225) (Table 1). There is substantial individual variability in the magnitude of changes in these cytokines in plasma after exercise (Figure 1).




Figure 1. Individual variability in plasma cytokine responses to the Western States 160 km Endurance Run. Data indicate the change (in pg/ml) from pre- to post-exercise. Data are combined from *Int J Sports Med* 24:541-547, 2003; *Brain Beh Immun* 19:398-403, 2005; *Brain Beh Immun* 20:578-584, 2006.

Table 1. Cytokine and chemokine expression in muscle and responses to mechanical strain and exercise.

	Expression in human muscle		Responsive to <i>in vivo</i> cyclic strain and EPS of C2C12 or human myotubes		Responsive to exercise		Change in plasma	
	mRNA	protein	mRNA	protein	mRNA	protein		
IL-1 β	✓	✓	✓	✓	✓		↑	↔
IL-1ra			✗	✗			↑↑↑	
IL-2		✓		✓				↓
IL-4	✓			✓				↔
IL-5				✓/✗				↔
IL-6	✓	✓	✓	✓	✓	✓	↑↑↑↑	
IL-7	✓	✓		✓/✗	✓			
IL-8	✓		✓	✓	✓		↑↑	↔
IL-10	✓			✗	✓		↑↑↑	
	(low)							
IL-12p35	✓			✗	✗		↑	
IL-13				✓				
IL-15	✓	✓		✓/✗	✓	✗	↑	↓
IL-18	✓	✓						
IFN- γ	✓	✓		✗				↔
TNF- α	✓	✓		✓/✗	✓/✗		↑	↔
MCP-1	✓	✓		✓	✓	✓	↑	
LIF	✓	✓		✓	✓	✗		↔
VEGF	✓	✓		✓/✗	✓	✓		
BDNF	✓	✓		✓	✗	✓	↑	↔
TGF- β	✓	✓			✓			↔
uPA	✓				✓			

✓ or ✗ indicate positive or negative evidence (respectively) of cytokine mRNA or protein expression/secretion in muscle and muscle contraction.

↑ or ↓ indicate an increase/decrease in plasma cytokine concentrations (more arrows denotes a greater increase). ↔ indicates no increase in plasma cytokine concentrations. (Inclusion of two symbols indicates conflicting data.)

N.B. Blank cells indicate no data are currently available. EPS, electromagnetic pulse stimulation. IFN, interferon; TNF, tumor necrosis factor; MCP, monocyte chemotactic protein; LIF, leukemia inhibitory factor; VEGF, vascular endothelial growth factor; BDNF, brain-derived neurotrophic factor; TGF, transforming growth factor; uPA, urokinase plasminogen activator.

Some of this variability appears to be related to variation in exercise intensity (156) and the extent of exercise-induced muscle damage (157, 228). By contrast, the plasma/serum concentrations of TNF- α (4 \times), brain-derived neurotrophic factor (BDNF) (2.5 \times), IL-1 β (2 \times), IL-12p40 (0.3 \times), IL-15 (0.05 \times) increase to a smaller extent, whereas leukemia inhibitory factor (LIF) and transforming growth factor (TGF)- β remain unchanged following exercise (25, 29, 39, 137, 153, 154, 186, 190, 220, 225).

The gene expression of IL-1 β , IL-6, IL-8, IL-10, IL-15, TNF- α , MCP-1, LIF and TGF- β in skeletal muscle increases following endurance exercise (39, 64, 68, 82, 102, 124, 147, 151, 152, 154, 208, 212) and resistance exercise (25, 50, 94, 124, 149, 150, 192) (Table 1). BDNF mRNA expression also tends to increase (non-significantly) in response to endurance exercise (137). Similar to plasma cytokine responses, exercise-induced changes in cytokine gene expression in muscle can also be highly variable, possibly due to variation in single nucleotide polymorphisms within cytokine genes (52, 186). However, less is known about changes in the protein abundance of these cytokines in skeletal muscle after exercise. IL-6 and BDNF protein expression increases in working skeletal muscle following endurance exercise (68, 89, 137), while the protein abundance of IL-6, IL-8 and MCP-1 also increases after resistance exercise (50, 51) and eccentric exercise (94). Changes in IL-1 β

are inconsistent (66, 129, 130), whereas IL-15 and LIF protein content does not change following exercise (25, 26, 130, 149).

In contrast with pro-inflammatory cytokines, exercise-induced changes in the anti-inflammatory cytokines IL-4 and IL-13 are less well characterized. We recently reported that IL-4 protein expression in muscle tended to decrease 2 h after resistance exercise in young men. However, following 12 weeks of resistance training, IL-4 protein expression increased when the same individuals performed another bout of resistance exercise (51). This finding suggests that IL-4 may play a role in muscular adaptations to training. In the same study, we observed that IL-13 protein expression in muscle did not change significantly after either bout of exercise (51).

CYTOKINE EXPRESSION AND SECRETION BY SKELETAL MUSCLE CELLS

Most of the evidence for cytokine expression in skeletal muscle is derived from the analysis of isolated RNA or protein extracts from homogenized muscle. However, muscle homogenates reflect contributions from intracellular, sequestered and interstitial sources of cytokines (23). This makes it difficult to identify specific cellular sources of cytokines *in vivo*. An extensive body of research has explored

Table 2. Expression of cytokines by skeletal muscle cells.

	Constitutive expression in myoblasts		Constitutive expression in myotubes		Change with differentiation	Stimulatory agents	Dose effect	Time effect
	mRNA	protein	mRNA	protein				
IL-1 β	H \checkmark (low) / *	H \checkmark		H \checkmark (low)	\uparrow (late)	TNF- α		
IL-1ra	C \checkmark			H *		LPS		+
IL-4	H *			H \checkmark (low)				
IL-6	H \checkmark / *	H \checkmark (low)	C \checkmark (low)	H \checkmark	\uparrow (early)	LPS	+	+
	C \checkmark (low)	C \checkmark		C \checkmark (low)		TNF- α	+	+
						IL-1 β	+	+
						IFN- γ	\leftrightarrow	\leftrightarrow
						TGF- β		
						HSP60	+	
						H ₂ O ₂	+	
						pyrogallol	+	
						X/XO	+	
						epinephrine	+	+
						MG-132		
						AP-1		
						HSF-1		
						Dithiothreitol		
						Thapsigargin		
						Tunicamycin	+	
						Castanospermine	+	
						ATP		+
IL-7			H \checkmark (low)	H \checkmark (low)	\uparrow	LPS		
IL-8	H \checkmark	H \checkmark (low)	H \checkmark	H \checkmark		TNF- α		
						IFN- γ		
						IL-1 β		
						'Hyper IL-6'		
						LPS		
						HSP60	+	
IL-10	H *			H \checkmark (low)		TNF- α		
IL-12				H \checkmark (low)		LPS		+
IL-13				H \checkmark (low)		IGF-1		
IL-15				H *	\uparrow	TNF- α		+
IL-17				H \checkmark (low)				
IL-18	C \checkmark							
TNF- α	H \checkmark / *	H *		H \checkmark (low)	\uparrow (early)	LPS	+	+
	C \checkmark (low)					TNF- α		
						IFN- γ		
MCP-1	H \checkmark / *	H *			\uparrow (early)	LPS		
						TNF- α		
						IFN- γ		
						IL-1 β		
						'Hyper IL-6'	+	+
						HSP60	+	
LIF			H \checkmark (low)	H \checkmark	\uparrow (early)	ionomycin		
IFN- γ	H \checkmark			H \checkmark				
BDNF			C \checkmark	C \checkmark	\downarrow			
VEGF-A				H \checkmark				
TGF- β	C \checkmark			H \checkmark	\uparrow	LPS		\leftrightarrow
						IL-1 α		+
						TNF- α		+
						IFN- γ		

\checkmark indicates constitutive expression. * indicates no constitutive expression. H, human muscle cells. C, C2C12 muscle cells. + indicates positive/additive response. \leftrightarrow indicates no effect. N.B. Blank cells indicate no data are currently available. MCP-1, monocyte chemotactic protein-1; LIF, leukemia inhibitory factor; IFN, interferon. BDNF, brain-derived neurotrophic factor; VEGF, vascular endothelial growth factor; TGF, transforming growth factor. LPS, lipopolysaccharide; HSP, heat shock protein; X/XO, xanthine/xanthine oxidase; hyper IL-6, IL-6 covalently linked to soluble IL-6 receptor; MG-132, proteasome inhibitor; AP-1, activator protein-1; HSF-1, heat shock factor 1;

the capacity of skeletal muscle cells to produce cytokines in vitro by culturing skeletal muscle cells with various agents, and by subjecting skeletal muscle cells to cyclic strain and electromagnetic pulse stimulation (Table 2).

Evidence from cell culture studies

C2C12 myoblasts, L6 muscle cells and human myoblasts constitutively express IL-8 (47, 133), IL-12 (72), IL-15 (180) and TGF- β (72). Some, but not all studies report that myoblasts

also constitutively express IL-1 β , IL-1ra, IL-6, TNF- α and MCP-1 (17, 47, 70-73, 106, 133, 144). Myoblasts do not express IL-10 or interferon (IFN)- γ (144). The constitutive expression of cytokines is generally stronger in differentiated myotubes compared with myoblasts (83, 85, 86, 102, 107, 134, 144, 235) (Table 2).

Myoblasts and myotubes express and secrete cytokines *in vitro* in response to a wide range of stimuli (Table 2). ATP, H₂O₂, xanthine/xanthine oxidase, nitric oxide and epinephrine are produced during exercise through redox reactions and autonomic nervous activity. These agents directly regulate the synthesis of cytokines by skeletal muscle cells by activating various signaling pathways (6, 70-72, 96, 107, 128, 235). The regulatory influence of TNF- α , IL-1 β , IL-6, IFN- γ , TGF- β and HSP60 is probably less direct, and depends on their relative concentration within the local microenvironment of skeletal muscle. Many of the agents listed above induce a concentration-dependent increase in the secretion of IL-6, IL-8, TNF- α and MCP-1 (70, 72, 73, 85, 107, 134, 144). LPS, TNF- α , IL-1 β , IL-6, IFN- γ and epinephrine also induce a time-dependent increase in the secretion of IL-6, TNF- α , TGF- β and MCP-1 for up to 48 hours (70-73, 133, 144). Human myotubes that undergo cyclic mechanical strain release IL-6, IL-8, MCP-1 and G-CSF (172). Mouse (62, 146, 235), rat (31) and human (25, 182, 195) myotubes that contract in response to electromagnetic pulse stimulation release a wide array of cytokines (Table 2). In addition to well characterized cytokines, the list of 'contraction-responsive' cytokines continues to grow (182, 195).

The findings from cell culture studies provide important information on skeletal muscle cell secretion of cytokines in the context of trauma, sepsis or chronic inflammatory conditions. However, these findings are not necessarily applicable to understanding how skeletal muscle cells generate cytokines during exercise, for several reasons. The concentrations of most of the agents are most likely lower in skeletal muscle during exercise compared with the concentrations used in cell culture studies. The period for which skeletal muscle is exposed to these agents during exercise is also typically shorter than the incubation periods used in cell culture studies. Although epinephrine stimulates IL-6 mRNA expression and secretion (70), epinephrine appears to play a relatively minor role in regulating systemic changes in IL-6 during exercise (214). Last, cyclic strain and electromagnetic pulse stimulation of myotubes do not fully reflect the dynamic conditions in skeletal muscle during exercise, where both stimulatory and inhibitory agents are present.

Evidence from in vivo muscle analysis

Human biopsy studies reveal that skeletal muscle expresses mRNA for numerous cytokines (Table 1) (25, 29, 83, 91, 94, 109, 149, 152, 154, 212). Mature myofibers make up most of the cellular mass within skeletal muscle, and as described in the previous section, myotubes express mRNA for various cytokines. Nevertheless, muscle homogenates represent a mixture of different types of cells; other inflammatory and stromal cells in skeletal muscle also secrete cytokines. Some research has used immunohistochemistry and immunofluorescence staining to examine the cellular sources of cytokines in

healthy human muscle at rest (68, 89, 149, 175). Malm et al. (129) found that IL-6 and IL-1 β are localized to both muscle and non-muscle cells. The same group subsequently reported that IL-6 expression is low within skeletal muscle cells themselves, whereas it is expressed in the epimysium (130). In muscle of patients with inflammatory myopathies, cytokines are highly abundant, but are mainly localized to other cell types such as inflammatory T cells and macrophages, and in proximity to blood vessels in the endomysium and perimysium (46, 194). Collectively, these findings demonstrate that cytokines are likely secreted by various cell types present in skeletal muscle, and not exclusively by skeletal muscle cells themselves.

Surprisingly little research has used RNA *in situ* hybridization, immunohistochemistry or immunofluorescence staining to identify which fibers and resident cell types (e.g., endothelial cells, infiltrating leucocytes, satellite cells) in skeletal muscle produce cytokines in response to exercise. Table 3 summarizes the results of studies that have attempted to identify where cytokines are expressed in skeletal muscle after exercise or muscle contractions. Hiscock et al. (89) report that following endurance exercise, both IL-6 mRNA and protein are expressed mainly in type II fibers with high muscle glycogen content. They propose that the greater IL-6 expression in type II fibers may result from greater release of Ca²⁺ from the sarcoplasmic reticulum into the cytosol of the type II fibers (89). Following eccentric exercise, Hubal et al. (94) observed that MCP-1 is mainly expressed within the interstitial space between myofibers, and localizes with macrophages and Pax7⁺ satellite cells. Two studies have demonstrated that IL-6 also localizes with Pax7⁺ satellite cells in the basal lamina of muscle fibers after eccentric exercise (138) and compensated hypertrophy (197). Following downhill running, Malm et al. (130) reported that IL-1 β is mainly localized to non-muscle cells, whereas Fielding et al. (66) found that IL-1 β is localized to the pericellular space. Using immunofluorescence staining for MCP-1 and IL-8, we recently reported that these chemokines are not present within mature myofibers after exercise. Rather, they are localized within the endomysium between muscle fibers and in close proximity to a number of cell types including macrophages, satellite cells and blood vessels (50). Hoier et al (90) investigated the subcellular localization of VEGF in muscle 2 h after cycling. They discovered that VEGF was located in the subsarcolemmal regions, between the contractile elements within the muscle fibers, and in pericytes positioned on the skeletal muscle capillaries. Lauritzen et al (116) conducted an elegant study in which they incorporated an enhanced green fluorescent protein tagged for IL-6 into muscle fibers isolated from the quadriceps muscles of mice. At rest, the fluorescent tag was localized in vesicular structures at the surface and in the interior of the transfected muscle fiber. Following *in situ* contractions of the fibers, the number of vesicles expressing the fluorescent tag decreased in both locations, indicating vesicular transport of IL-6 out of the fiber. These diverse findings highlight the need for more research to gain a better understanding of the local regulation and secretion of cytokines in muscle during exercise.

Table 3. Cytokine localization in skeletal muscle after exercise.

Cytokine	Species	Exercise mode	Muscle	Located within muscle fibers	Located outside muscle fibers	Ref.
IL-6	Humans	Endurance	Vastus lateralis	✓ (type II fibers)		(89)
IL-6*	Rats	Endurance	Plantaris	✓ (type I and IIa fibers)		
IL-6	Humans	Eccentric	Vastus lateralis		✓ (satellite cells)	(138)
IL-6	Humans	Eccentric	Vastus lateralis	low	✓ (fibroblasts)	(130)
IL-6	Humans	Eccentric	Vastus lateralis	low	low	(129)
IL-6	Mice	Eccentric	Gastroc.	✓	✓ (inflammatory and satellite cells)	(226)
LIF	Humans	Eccentric	Vastus lateralis		✓ (endothelial cells)	(130)
IL-1 α	Humans	Eccentric	Vastus lateralis	✓	✓ (endothelial cells)	(129)
IL-1 β	Humans	Eccentric	Vastus lateralis	✓	✓	(129)
IL-1 β	Humans	Eccentric	Vastus lateralis		✓ (pericellular space)	(66)
MCP-1	Humans	Eccentric	Vastus lateralis		✓ (macrophages and satellite cells)	(94)
MCP-1	Humans	Resistance	Vastus lateralis		✓ (macrophages, satellite cells, blood vessels)	(50)
IL-8	Humans	Resistance	Vastus lateralis		✓ (macrophages, blood vessels)	(50)
VEGF	Humans	Endurance	Vastus lateralis	✓ (subsarcolemmal sarcoplasm)	✓ (pericytes)	(90)
VEGF*	Rats	Endurance	Plantaris	✓ (type IIb fibers)		(22)
VEGF*	Rats	Endurance	Gastroc.	✓ (type I and IIa fibers)		(28)
TGF- β ₁	Rats	Eccentric	Gastroc.	✓ (injured myofibers)		(201)

* mRNA. Gastroc, gastrocnemius.

DISSOCIATION BETWEEN LOCAL AND SYSTEMIC CYTOKINE RESPONSES TO EXERCISE

Curiously, although many cytokines are expressed in skeletal muscle following exercise, with the exception of IL-6, they are not released into the circulation—at least in large amounts (65, 68, 211, 212). In explanation of these observations, it has been suggested that some cytokines are produced locally by interstitial cells, and may not enter the circulation (130, 191). Catoire et al (37) recently conducted a systematic comparison of cytokine gene expression in muscle and plasma cytokine concentrations after one-legged 1 h cycling at 60% heart rate reserve. They discovered that mRNA expression of IL-6, MCP-1, CXCL2 (macrophage inflammatory protein-2 α) and CX3CL1 (fractalkine) was significantly upregulated. By contrast, mRNA expression of other cytokines including IL-7, IL-8, IL-15 and BDNF did not change significantly after exercise. Within plasma, MCP-1 and fractalkine increased after exercise, whereas IL-6 remained unchanged. This is the only study to report simultaneous changes in MCP-1 and fractalkine within muscle and plasma following exercise. The finding that plasma IL-6 concentration did not change despite local expression in muscle may reflect the relatively small muscle mass and low intensity of exercise (37). An alternative explanation for the dissociation between local and systemic cytokine responses is that skeletal muscle may not secrete sufficient quantities of cytokines to increase their concentration in the systemic circulation.

Although distinct from exercise, Borge et al. (23) conducted an elegant study to investigate whether cytokines are released systemically from skeletal muscle in response to lipopolysaccharide (LPS). Plasma and interstitial fluid were collected from mice 0.5, 1.5 and 3 h after intravenous administration of 3.5 mg/kg LPS. The concentrations and kinetics of changes in cytokines were markedly different between interstitial fluid and plasma. The findings from this study by Borge et al. (23) provide important information about skeletal muscle as a source of circulating cytokines. The higher concentration of IL-1 β in interstitial fluid compared with plasma suggests that although skeletal muscle cells produce IL-1 β , the systemic release of IL-1 β from skeletal muscle is probably tightly regulated. The higher concentrations of TNF- α , IL-10, MCP-1 in plasma compared with interstitial fluid 1.5 h after LPS infusion suggests that skeletal muscle is not a major source of these cytokines in the circulation. The smaller difference in the concentrations of IL-6 in plasma and interstitial fluid is consistent with other evidence that skeletal muscle releases IL-6 into the circulation during exercise (68, 211, 212). The time course of changes in the secretion of cytokines in this study is similar to that reported in plasma following exercise (160). The early secretion of TNF- α and IL-1 β probably stimulated the sustained production of IL-6 (44, 71, 72, 125, 162). Constitutive expression of IL-10 in skeletal muscle cells is low (144), but IL-10 production may have increased in response to the early rise in TNF- α and IL-1 β secretion (67).

OTHER SOURCES OF LOCAL AND SYSTEMIC CYTOKINES DURING EXERCISE

Cells within the microvasculature, namely endothelial cells and pericytes, are important regulators of angiogenesis and myogenesis, making them key players in both muscle and vascular generation following injury (2). Endothelial cells (78, 100, 111, 217, 238) and pericytes secrete various cytokines (41, 108). Fibroblasts contribute to production of the extracellular matrix of muscle connective tissue by secreting fibronectin, laminin, specific tenascins and neural cell adhesion molecules (141). In response to muscle injury, fibroblasts proliferate and begin to produce collagen-rich extracellular matrix to restore the muscle’s framework (121). Fibroblasts also secrete assorted cytokines (47, 78, 118, 136, 163). Neutrophils play an important role in breaking down damaged muscle tissue in the acute phase of muscle injury (148, 174), whereas monocytes/macrophages regulate subsequent tissue regeneration (12, 218). Neutrophils (36, 58, 132, 187, 223) and monocytes/macrophages (8, 48, 54, 78, 93, 123, 131, 179) both secrete a variety of cytokines. As cytokine-producing cells, endothelial cells, pericytes, fibroblasts, neutrophils and monocytes/macrophages may all contribute to global cytokine expression in skeletal muscle.

In addition to skeletal muscle, IL-6 is also released from the brain (158) and peritendinous tissue (114) after exercise. Whereas IL-6 mRNA is expressed in adipose tissue following exercise (45, 103), IL-6 is not released from adipose tissue during exercise (126). Evidence indicates that macrophages secrete IL-1β in the brain following downhill running (35), but it is unknown whether IL-1β is released from the brain into the circulation during exercise. Exercise also increases gene expression of IL-1ra and IL-15 receptorα in the liver in fasting rats (30), but it remains unknown if the liver is a source of circulating cytokines after exercise.

Leucocytes are probably only a minor source of circulating cytokines following exercise (Table 4). Studies on cytokine production by leucocytes can be divided into those that have measured cytokine gene expression in leucocytes, intracellular cytokine production, and extracellular cytokine secretion. Leucocyte mRNA expression of IL-1β, IL-1ra, IL-8 and IL-10 increases, whereas IL-6 mRNA expression remains

unchanged after exercise (1, 20, 154, 161). Monocyte intracellular cytokine production of IL-1β, IL-6, TNF-α, BDNF increases, decreases or remains unchanged following exercise (27, 185, 207, 209, 210). Extracellular cytokine secretion by mononuclear cells or in whole blood stimulated with LPS is also variable (1, 18, 19, 56, 57, 81, 87, 112, 173, 189, 202, 231). Fewer studies have investigated changes in cytokine expression or secretion by T lymphocytes following exercise. Two studies have reported that the number of IFN-γ+ T cells decreases after exercise, while the number of IL-4+ T cells remains unchanged (113, 213). In contrast with these findings, Zaldivar et al (239) found that the percentage of T cells that expressed IL-4, IL-6 and TNF-α increased following exercise. Kakani et al (99) also observed that the secretion of both Th1 cytokines (IL-2 and TNF-α) and Th2 cytokines (IL-6, IL-10) by T cells stimulated with phytohemagglutinin increased after exercise. Work by La Voy et al (117) demonstrated that cytokine production during exercise may depend on changes in the numbers certain subsets of T cells.

Skeletal muscle has been proposed as the dominant source of circulating IL-6 based on the increase in the arterial-femoral venous differences in the concentration of IL-6 (215). However, this does not provide *prima facie* evidence that skeletal muscle cells are the main cell type that secretes cytokines such as IL-6 into the circulation during exercise. Skeletal muscle is composed of many other cell types such as fibroblasts, myeloid cells, pericytes which also secrete cytokines. The organs and cells that secrete other cytokines (not produced in abundance by skeletal muscle cells) into the circulation during exercise remain to be determined. The results from many of these studies on cytokine secretion by leucocytes *in vitro* highlight the importance of how we interpret such data. It is important to consider differences in stimulated versus spontaneous cytokine secretion, intracellular cytokine production versus extracellular secretion, and changes in the absolute amount of cytokines that are secreted versus cytokine secretion per cell.

REGULATION OF CYTOKINE SYNTHESIS AND SECRETION

Cytokine secretion by skeletal muscle cells involves various intracellular factors, including mitogen-activated protein

Table 4. Leucocyte cytokine mRNA expression and secretion in response to exercise.

	mRNA expression	Unstimulated production	Stimulated production	Production per cell	Change in plasma
IL-1β	↑	↓	↓	↓	↔
IL-1ra	↑	↑	↔	↔	↑↑↑
IL-6	↓	↔	↑	↑ ↓ ↔	↑↑↑↑
IL-8	↑				↑↑ ↔
IL-10	↑	↓	↓	↓	↑↑↑
TNF-α		↓	↓	↔ ↑ ↓ ↔	↑ ↔

↑ or ↓ indicate an increase/decrease in plasma cytokine concentrations (more arrows denotes a greater increase). ↔ indicates no increase in plasma cytokine concentrations. (Inclusion of two symbols indicates conflicting data.). Unstimulated production refers to assays in which whole blood or cells were incubated with no external agent. Stimulated production refers to assays in which whole blood or cells were incubated with an external agent such as lipopolysaccharide.

kinases, heat shock factor 1, histone deacetylases and transcription factors such as nuclear factor of activated T cells (NFAT), activating protein (AP)-1 and NF κ B (6, 70-72, 107, 128, 233, 212, 235). In addition, cellular processes such as Ca²⁺ signaling and protein unfolding also stimulate muscle cells to express cytokine genes and/or secrete cytokines (92, 233). Activating transcription factor 3 is an important regulator of cytokine secretion by macrophages (76), and may also play a role in skeletal muscle cells. In comparison with our knowledge of the factors that induce cytokine synthesis, much less is known about the factors that restrict and/or inhibit cytokine expression and secretion (145). In the context of exercise, this information is important because it could account for why IL-1 β and TNF- α mRNA expression in skeletal muscle increases, yet the circulating concentrations of these cytokines remains comparatively low following exercise (150-152, 212). Below, we propose some potential negative regulatory mechanisms that may govern cytokine expression and secretion by skeletal muscle cells.

RNA-binding proteins

Intracellular utilization of mRNA depends on several processes including mRNA maturation, shuttling and stability. In turn, these post-transcriptional processes are under the control of RNA-binding proteins and microRNAs. RNA-binding proteins regulate mRNA utilization by binding to adenine/uracil-rich elements downstream of the 3' untranslated regions of transcripts (205). A small subset of RNA-binding proteins are active in skeletal muscle, including human antigen R, KH-type splicing regulatory protein, CUG binding protein 1, poly(A) binding protein, Lin-28 and tristetraprolin (9). These RNA-binding proteins are known to regulate myogenesis (9), but they may also control cytokine translation in skeletal muscle cells. Human antigen R enhances the stability of TNF- α mRNA (49, 140). However, together with CUG binding protein 1 and tristetraprolin, human antigen R may also silence TNF- α translation (33, 98, 101, 240). Tristetraprolin expression increases (221), whereas the expression of human antigen R decreases (181) in response to the anti-inflammatory cytokines IL-4 and IL-10.

Relatively little research has examined changes in the expression of RNA-binding proteins in skeletal muscle following exercise. Hubal et al. (94) reported that mRNA expression of zinc finger protein 36 (a member of the tristetraprolin family) increases in skeletal muscle following exercise. This response is greater following eccentric exercise compared with concentric exercise, and is augmented following repeated bouts of eccentric exercise (94) which suggests a role for zinc finger protein 36 in regulating skeletal muscle adaptation following injury. Another study observed that tristetraprolin mRNA expression precedes that of LPS-inducible CXC chemokine mRNA expression in C2C12 myoblasts and in skeletal muscle following freeze injury (193). Geyer et al (75) performed an elegant study in which they induced the expression of the RNA-binding protein tristetraprolin in C2C12 myotubes. This treatment suppressed mRNA expression of MCP-1, KC (IL-8) and IL-6, while it also reduced MCP-1 secretion following LPS stimulation. These findings highlight the need for further research to gain greater insights into the role of RNA-binding proteins in regulating cytokine secretion by skeletal muscle cells.

MicroRNAs

MicroRNAs also bind to the 3' untranslated regions of transcripts (222) and interact with RNA-binding proteins to regulate the fate of mRNA (59, 98, 127). MicroRNAs such as Let-7, miR-146, miR-221, miR-155 and miR-106 regulate the expression of IL-1, IL-6, IL-8, TNF- α and IL-10 by immune cells (21, 38, 59, 98, 170, 198, 224, 229). Skeletal muscle cells express a number of microRNAs, including miR-133, miR-1, miR-367, miR-135a, miR-222, miR-29a, b and c, miR-221, miR-223 and miR-206 (34, 42, 105). MicroRNA Let-7 inhibits the secretion of IL-13 in human myotubes (97). Other microRNAs such as miR-367, miR-222, and miR-29 may control cytokine secretion by skeletal muscle cells indirectly by altering the activity of endothelial nitric oxide synthase (188) and signal transducer and activator of transcription (STAT) proteins (53). As more microRNAs are identified in skeletal muscle, this may improve our knowledge of whether they regulate cytokine expression and secretion by skeletal muscle cells.

Suppressor of cytokine signaling (SOCS)

Surprisingly little research has investigated the role of SOCS proteins in regulating cytokine synthesis and signaling by skeletal muscle cells. Paradoxically, the limited evidence available indicates that overexpression of SOCS3 increases IL-6 transcription in myotubes (204). Under some conditions, the interaction between SOCS proteins and cytokines may be reciprocal. IL-6 and TNF- α induce SOCS3 mRNA expression in C2 myoblasts (5) and cardiac myoblasts (230). TNF- α infusion *in vivo* also stimulates SOCS3 mRNA in murine muscle (60). Further research is warranted to examine in greater detail the function of SOCS proteins as regulators of cytokine synthesis and signaling by skeletal muscle cells.

Soluble receptors

Soluble receptors can also restrict cytokine signaling through two main mechanisms. First, soluble receptors can act as a 'non-signaling sink' that directly competes with membrane-bound receptors for ligand binding. If the ligand-binding affinity of soluble receptors and membrane-bound receptors is similar, the capacity of soluble receptors to inhibit signaling depends on the balance between the two types of receptors (84). Soluble receptors for IL-1 and TNF- α appear to operate in this manner (3, 10). Second, soluble receptors can arise through the proteolytic cleavage of membrane-bound receptors. This process results in fewer membrane-bound receptors to bind ligands and initiate cell signaling (84). TNF- α and IL-6 can also induce shedding and/or endocytosis of their own receptors (55, 79, 88, 177). These actions may represent an autocrine negative feedback loop to prevent excess ligand stimulation. Currently, there is insufficient evidence to determine whether soluble receptors regulate cytokine signaling in skeletal muscle following exercise. Gene expression of soluble IL-6 receptor (but not gp130) is elevated in skeletal muscle between 4.5 and 9 h after exercise (104), whereas the plasma concentration of soluble IL-6 receptor does not change (104) or only increases slightly (120). The presence of circulating soluble IL-6 receptors after exercise may depend on proteolytic cleavage of IL-6 receptors (120). Gene expression of TNF- α and IL-1 β in skeletal muscle is elevated for up to 24 h after exercise (124, 150-152). Although the plasma concen-

trations of soluble TNF- α receptors and IL-1ra are also elevated for several hours after exercise (152, 160, 225), it is unknown whether these receptors are derived from, and are active in skeletal muscle.

Other factors

Other factors such as IL-6 and HSP72 produced locally in skeletal muscle during exercise may also regulate cytokine synthesis. IL-6 inhibits LPS-induced synthesis of TNF- α by monocytes (196). In response to LPS treatment *in vivo*, TNF- α concentration is lower in serum and broncho-alveolar lavage fluid from IL-6^{+/+} mice compared with IL-6^{-/-} mice (236). An increase in plasma IL-6 concentration following IL-6 infusion or exercise inhibits the systemic release of TNF- α in response to LPS (206). Local production of IL-6 may therefore regulate the synthesis and systemic release of TNF- α during exercise.

In addition to cytokines, endurance exercise induces HSP72 mRNA expression in skeletal muscle (16). Heat exposure increases HSP72 mRNA expression and IL-6 mRNA expression and protein synthesis in C2C12 myotubes in a temperature-dependent manner (234). As further evidence for this regulatory role of HSP72, the heat shock inhibitor Knk437 attenuates HSP72 mRNA expression, and completely blocks IL-6 mRNA expression in myotubes incubated at 42°C (234). The interaction between HSP72 and IL-6 appears to be reciprocal, because IL-6 infusion induces HSP72 mRNA expression in skeletal muscle (63). Huey and Meador (95) demonstrated that IL-6 regulates the expression of HSPs in skeletal muscle in response to LPS, but not exercise. In direct contrast with IL-6, heat exposure inhibits TNF- α mRNA expression in myotubes (234) and TNF- α protein synthesis in other cell types (61, 142). These findings implicate HSP72 as a negative regulator of TNF- α mRNA expression and synthesis, but further research is required to confirm this notion. The upstream regulator of HSPs, heat shock factor -1 may play a more central role than HSPs in regulating cytokine secretion by skeletal muscle cells (233). Ohno et al (159) have also demonstrated that acute heat stress suppresses NF κ B activity in C2C12 muscle cells. This response was accompanied by increased expression of HSP72 (159), but it remains to be determined if HSP72 can block NF κ B activity in muscle cells.

Skeletal muscle expresses the gene for tumor necrosis factor receptor-associated factor (TRAF)-6-inhibitory zinc finger protein (TIZ) (77). By inhibiting the activation of NF κ B, c-Jun N-terminal kinase and AP-1 (199), zinc finger proteins such as TIZ may restrict cytokine secretion by skeletal muscle cells. Similar to other factors described above, future research could investigate the regulatory roles of zinc finger proteins in skeletal muscle.

In another interesting study, Lee (119) demonstrated that treatment of C2C12 myotubes with IGF-1 inhibited mRNA expression of IL-6 and TNF- α . This effect was due to suppression of TLR4 signaling, which was in turn mediated by inhibition of the PI3K/Akt signaling pathway (119). These findings provide further evidence of potential autocrine loops and cross-talk between cytokines and growth factors within skeletal muscle.

Cytokine trafficking and secretion

In contrast with cells of the immune system, regulation of cytokine trafficking and secretion in muscle cells remains largely unknown. Hoier et al (90) made the first attempt to characterize the subcellular localization of VEGF in skeletal muscle. However, they did not investigate dynamic changes in trafficking of VEGF within muscle cells. Lauritzen et al (116) have provided the most detailed insights to date on the mechanisms of cytokine secretion by skeletal muscle fibers. Stow and Murray (216) have provided a comprehensive overview on the mechanisms of trafficking and secretion in immune cells, and this may be used as a guide to new research in muscle cells.

REGULATION OF CYTOKINE EXPRESSION AND SECRETION IN MYOBLASTS VERSUS MYOTUBES

As noted previously, and summarized in Table 2, the constitutive expression of cytokines varies between myoblasts and myotubes. Below we discuss some of these differences, some of the mechanisms that may govern alterations in cytokine expression and secretion by skeletal muscle cells as they differentiate, and the possible biological significance of these differences.

Gene expression of numerous chemokines and their receptors increases markedly after 16–48 h of differentiation (80). Of note, MCP-1 mRNA expression peaks at 16 h, whereas TNF- α mRNA expression increases 10-fold between 16 and 24 h and declines thereafter (80). Several studies indicate that myoblasts and myotubes constitutively express similar levels of IL-6 mRNA. However, compared with myotubes, myoblasts produce substantially more IL-6 protein upon stimulation with IL-1 β , TNF- α and LPS (71, 144, 178). In response to TNF- α and LPS, myoblast production of IL-6 increases in a linear manner, whereas myotube production of IL-6 increases in a more ‘bell-shaped’ manner (178). The greater sensitivity of myoblasts to pro-inflammatory stimuli such as TNF- α and LPS may reflect the requirement for myoblasts to secrete factors such as IL-6 to promote myoblast proliferation (13, 110), and therefore, muscle regeneration (178). Protein expression of MCP-1 (85), IL-6 (15) and LIF (25) also increases during the first 24–48 h of differentiation. Protein expression of IL-7 and IL-15 increases more steadily between 2–7 d of differentiation (83, 180). Protein expression of IL-1 β and IL-1ra (14) and MCP-1 secretion (40) increase after 12–16 d of differentiation. Gene and protein expression BDNF expression decreases after 4 d (143).

The factors responsible for changes in cytokine expression by skeletal muscle cells as they differentiate are uncertain. Some of the factors that regulate cytokine expression in skeletal muscle cells (e.g., mitogen-activated protein kinases, histone deacetylases, NFAT and NF κ B) also control muscle cell differentiation (11, 15, 139, 169), and may therefore account for alterations in cytokine expression and secretion in skeletal muscle cells as they differentiate. The expression of some toll-like receptors (e.g., TLR2, TLR5) increases during muscle cell differentiation (24), which may also influence signaling path-

ways linked to cytokine expression and secretion in skeletal muscle cells (69). Changes in cytokine expression during muscle cell differentiation appear to play an important functional role. For example, increased expression of IL-6 (15) and reduced expression of BDNF (143) in skeletal muscle cells is necessary for them to differentiate. Conversely, increased expression of TNF- α during muscle cell differentiation (80) may inhibit myoblast differentiation (115, 227), although this effect is not entirely consistent (43). Using an RNAi screen, Ge et al (74) identified more than 100 cytokines that act regulate myoblast differentiation. Based on their results, they were able to classify these cytokines according to their capacity to initiate differentiation, regulate myocyte fusion and inhibit differentiation. Increased chemokine expression during differentiation may also control the migration and/or positioning of myoblasts so that they can successfully fuse with nascent myotubes (80). Alternatively, increased cytokine expression during muscle cell differentiation may promote muscle cell proliferation (122, 237), and migration of monocytes (40) and mesenchymal stem cells (176) to support muscle growth. Last, autocrine cross-talk between cytokines may also control muscle cell differentiation (4, 7, 71, 125).

CONCLUSION

Our understanding of the importance of skeletal muscle and cytokines as mediators of metabolism has increased substantially over the last decade or so. Research to date has identified more than 600 different proteins that are secreted by skeletal muscle cells (168). In this rapidly advancing age of 'omics' technologies, the muscle cell secretome will continue to grow and provide new targets on which to focus. We now know that myokines exert various endocrine effects on various metabolically active organs, including adipose tissue, the liver, the pancreas and the brain. Nevertheless, perhaps with the exception of IL-6 and LIF, our knowledge of the factors and mechanisms that regulate cytokine production and release from skeletal muscle cells during exercise remains somewhat limited. Skeletal muscle cells produce numerous cytokines in response to various agents, but not all these agents are present at similar concentrations and/or are active in the muscle microenvironment during exercise. The capacity for these individual agents to stimulate skeletal muscle cells to produce cytokines is well characterized. However, the control of cytokine production in skeletal muscle during exercise is more complex than the *in vitro* setting, and depends on interactions between a variety of local and systemic factors. Future research should aim to treat skeletal muscle cells with combinations and concentrations of agents that are present in skeletal muscle during exercise.

Cyclic strain and electromagnetic stimulation of skeletal muscle myotubes *in vitro* has generated useful insights into the signaling pathways that govern cytokine production by skeletal muscle cells during exercise. Adding other factors—such as cytokines themselves—to this experimental system could simulate the muscle microenvironment during exercise. In doing so, this approach may assist in characterizing interactions between factors that stimulate or inhibit the ability of skeletal muscle cells to produce cytokines during exercise.

More research is warranted to identify the feedback mechanisms that govern cytokine synthesis by skeletal muscle cells; in particular, which mechanisms are most important, how they interact with each other, and how they are induced and regulated. These research endeavors are important for several reasons. First, this information may help to understand the factors governing the systemic release of cytokines. Second, this information may help to understand the processes that regulate acute inflammatory responses to tissue injury. Last, this information may help to determine why some pro-inflammatory cytokines are chronically elevated in skeletal muscle of patients with idiopathic myopathies, rheumatoid arthritis and muscular dystrophy.

Finally, much has been written about the anti-inflammatory effects of exercise. Petersen et al (171) first proposed the notion that exercise-induced increases in IL-6, IL-1ra and IL-10 exert beneficial anti-inflammatory effects to counteract obesity and insulin resistance. Although this theory is appealing from a mechanistic perspective (206), cytokines may play a relatively minor role in regulating the health benefits of exercise training. Evidence in support of this notion is that brisk walking does not stimulate any discernible increase in circulating cytokines (135), yet regular walking is associated with many health benefits. Furthermore, marathon running induces high physiological stress and a large cytokine response (155), but it is doubtful the cytokines are increased for the purposes of health. Instead, exercise-induced cytokine changes may represent a more generalized response to internal and/or external stress. Factors such as oxidative or nitrosative stress, damaged or unfolded proteins, hyperthermia or energy imbalance likely induce cytokine production during exercise through catecholamines, endotoxin, alarmins, ATP and pro-inflammatory cytokines themselves (232). These issues highlight the need for further research to enhance our understanding of the biological significance of exercise-induced cytokine responses.

REFERENCES

1. Abbasi A, Fehrenbach E, Hauth M, Walter M, Hudemann J, Wank V, Niess AM, and Northoff H. Changes in spontaneous and LPS-induced *ex vivo* cytokine production and mRNA expression in male and female athletes following prolonged exhaustive exercise. *Exerc Immunol Rev* 19: 8-28, 2013.
2. Abou-Khalil R, Mounier R, and Chazaud B. Regulation of myogenic stem cell behavior by vessel cells: the "menage a trois" of satellite cells, periendothelial cells and endothelial cells. *Cell Cycle* 9: 892-896, 2010.
3. Aderka D, Engelmann H, Maor Y, Brakebusch C, and Wallach D. Stabilization of the bioactivity of tumor necrosis factor by its soluble receptors. *J Exp Med* 175: 323-329, 1992.
4. Al-Shanti N, Saini A, Faulkner S, and Stewart C. Beneficial synergistic interactions of TNF- α and IL-6 in C2 skeletal myoblasts--potential cross-talk with IGF system. *Growth Factors* 26: 61-73, 2008.
5. Al-Shanti N, and Stewart CE. Inhibitory effects of IL-6 on IGF-1 activity in skeletal myoblasts could be mediated by the activation of SOCS-3. *J Cell Biochem* 113: 923-933, 2012.

6. Allen DL, Uyenishi JJ, Cleary AS, Mehan RS, Lindsay SF, and Reed JM. Calcineurin activates interleukin-6 transcription in mouse skeletal muscle in vivo and in C2C12 myotubes in vitro. *Am J Physiol Regul Integr Comp Physiol* 298: R198-210, 2010.
7. Alvarez B, Quinn LS, Busquets S, Quiles MT, Lopez-Soriano FJ, and Argiles JM. Tumor necrosis factor- α exerts interleukin-6-dependent and -independent effects on cultured skeletal muscle cells. *Biochim Biophys Acta* 1542: 66-72, 2002.
8. Anegón I, Moreau JF, Godard A, Jacques Y, Peyrat MA, Hallet MM, Wong G, and Soullillou JP. Production of human interleukin for DA cells (HILDA)/leukemia inhibitory factor (LIF) by activated monocytes. *Cell Immunol* 130: 50-65, 1990.
9. Apponi LH, Corbett AH, and Pavlath GK. RNA-binding proteins and gene regulation in myogenesis. *Trends Pharmacol Sci* 32: 652-658, 2011.
10. Arend WP, Malyak M, Smith MF, Jr., Whisenand TD, Slack JL, Sims JE, Giri JG, and Dower SK. Binding of IL-1 α , IL-1 β , and IL-1 receptor antagonist by soluble IL-1 receptors and levels of soluble IL-1 receptors in synovial fluids. *J Immunol* 153: 4766-4774, 1994.
11. Armand AS, Bourajjaj M, Martinez-Martinez S, el Azzouzi H, da Costa Martins PA, Hatzis P, Seidler T, Redondo JM, and De Windt LJ. Cooperative synergy between NFAT and MyoD regulates myogenin expression and myogenesis. *J Biol Chem* 283: 29004-29010, 2008.
12. Arnold L, Henry A, Poron F, Baba-Amer Y, van Rooijen N, Plonquet A, Gherardi RK, and Chazaud B. Inflammatory monocytes recruited after skeletal muscle injury switch into antiinflammatory macrophages to support myogenesis. *J Exp Med* 204: 1057-1069, 2007.
13. Austin L, and Burgess AW. Stimulation of myoblast proliferation in culture by leukaemia inhibitory factor and other cytokines. *J Neurol Sci* 101: 193-197, 1991.
14. Authier FJ, Chazaud B, Plonquet A, Eliezer-Vanerot MC, Poron F, Belec L, Barlovatz-Meimon G, and Gherardi RK. Differential expression of the IL-1 system components during in vitro myogenesis: implication of IL-1 β in induction of myogenic cell apoptosis. *Cell Death Differ* 6: 1012-1021, 1999.
15. Baeza-Raja B, and Munoz-Canoves P. p38 MAPK-induced nuclear factor- κ B activity is required for skeletal muscle differentiation: role of interleukin-6. *Mol Biol Cell* 15: 2013-2026, 2004.
16. Bartlett JD, Hwa Joo C, Jeong TS, Louhelainen J, Cochran AJ, Gibala MJ, Gregson W, Close GL, Drust B, and Morton JP. Matched work high-intensity interval and continuous running induce similar increases in PGC-1 α mRNA, AMPK, p38, and p53 phosphorylation in human skeletal muscle. *J Appl Physiol* 112: 1135-1143, 2012.
17. Bartoccioni E, Michaelis D, and Hohlfeld R. Constitutive and cytokine-induced production of interleukin-6 by human myoblasts. *Immunol Lett* 42: 135-138, 1994.
18. Bassit RA, Sawada LA, Bacurau RF, Navarro F, Martins E, Jr., Santos RV, Caperuto EC, Rogeri P, and Costa Rosa LF. Branched-chain amino acid supplementation and the immune response of long-distance athletes. *Nutrition* 18: 376-379, 2002.
19. Baum M, Muller-Steinhardt M, Liesen H, and Kirchner H. Moderate and exhaustive endurance exercise influences the interferon- γ levels in whole-blood culture supernatants. *Eur J Appl Physiol* 76: 165-169, 1997.
20. Bernecker C, Scherr J, Schinner S, Braun S, Scherbaum WA, and Halle M. Evidence for an exercise induced increase of TNF- α and IL-6 in marathon runners. *Scand J Med Sci Sports* 23: 207-214, 2013.
21. Bhaumik D, Scott GK, Schokrpur S, Patil CK, Orjalo AV, Rodier F, Lithgow GJ, and Campisi J. MicroRNAs miR-146a/b negatively modulate the senescence-associated inflammatory mediators IL-6 and IL-8. *Aging* 1: 402-411, 2009.
22. Birot OJ, Koulmann N, Peinnequin A, and Bigard XA. Exercise-induced expression of vascular endothelial growth factor mRNA in rat skeletal muscle is dependent on fibre type. *J Physiol* 552: 213-221, 2003.
23. Borge BA, Kalland KH, Olsen S, Bletsa A, Berggreen E, and Wiig H. Cytokines are produced locally by myocytes in rat skeletal muscle during endotoxemia. *Am J Physiol Heart Circ Physiol* 296: H735-744, 2009.
24. Boyd JH, Divangahi M, Yahiaoui L, Gvozdic D, Qureshi S, and Petrof BJ. Toll-like receptors differentially regulate CC and CXC chemokines in skeletal muscle via NF- κ B and calcineurin. *Infect Immun* 74: 6829-6838, 2006.
25. Broholm C, Laye M, Brandt C, Vadalasetty R, Pilegaard H, Pedersen BK, and Scheele C. LIF is a contraction-induced myokine stimulating human myocyte proliferation. *J Appl Physiol* 111: 251-259, 2011.
26. Broholm C, Mortensen OH, Nielsen S, Akerstrom T, Zankari A, Dahl B, and Pedersen BK. Exercise induces expression of leukaemia inhibitory factor in human skeletal muscle. *J Physiol* 586: 2195-2201, 2008.
27. Brunelli A, Dimauro I, Sgro P, Emerenziani GP, Magi F, Baldari C, Guidetti L, Luigi LD, Parisi P, and Caporossi D. Acute exercise modulates BDNF and pro-BDNF protein content in immune cells. *Med Sci Sports Exerc* 44: 1871-1880, 2012.
28. Brutsaert TD, Gavin TP, Fu Z, Breen EC, Tang K, Mathieu-Costello O, and Wagner PD. Regional differences in expression of VEGF mRNA in rat gastrocnemius following 1 hr exercise or electrical stimulation. *BMC Physiol* 2: 8, 2002.
29. Buford TW, Cooke MB, and Willoughby DS. Resistance exercise-induced changes of inflammatory gene expression within human skeletal muscle. *Eur J Appl Physiol* 107: 463-471, 2009.
30. Buler M, Aatsinki SM, Skoumal R, Komka Z, Toth M, Kerkela R, Georgiadi A, Kersten S, and Hakkola J. Energy-sensing factors coactivator peroxisome proliferator-activated receptor γ coactivator 1- α (PGC-1 α) and AMP-activated protein kinase control expression of inflammatory mediators in liver: induction of interleukin 1 receptor antagonist. *J Biol Chem* 287: 1847-1860, 2012.
31. Bustamante M, Fernandez-Verdejo R, Jaimovich E, and Buvinic S. Electrical stimulation induces IL-6 in skeletal muscle through extracellular ATP by activating Ca²⁺ signals and an IL-6 autocrine loop. *Am J Physiol Endocrinol Metab* 306: E869-882, 2014.
32. Cannon JG, and Kluger MJ. Endogenous pyrogen activity in human plasma after exercise. *Science* 220: 617-619, 1983.
33. Carballo E, Lai WS, and Blackshear PJ. Feedback inhibition of macrophage tumor necrosis factor- α production by tristetraprolin. *Science* 281: 1001-1005, 1998.
34. Cardinali B, Castellani L, Fasanaro P, Basso A, Alema S, Martelli F, and Falcone G. MicroRNA-221 and microRNA-222 modulate differentiation and maturation of skeletal muscle cells. *PLoS One* 4: e7607, 2009.

35. Carmichael MD, Davis JM, Murphy EA, Carson JA, Van Rooijen N, Mayer E, and Ghaffar A. Role of brain macrophages on IL-1 β and fatigue following eccentric exercise-induced muscle damage. *Brain Behav Immun* 24: 564-568, 2010.
36. Cassatella MA, Bazzoni F, Ceska M, Ferro I, Baggioolini M, and Berton G. IL-8 production by human polymorphonuclear leukocytes. The chemoattractant formyl-methionyl-leucyl-phenylalanine induces the gene expression and release of IL-8 through a pertussis toxin-sensitive pathway. *J Immunol* 148: 3216-3220, 1992.
37. Catoire M, Mensink M, Kalkhoven E, Schrauwen P, and Kersten S. Identification of human exercise-induced myokines using secretome analysis. *Physiol Genomics* 46: 256-267, 2014.
38. Ceppi M, Pereira PM, Dunand-Sauthier I, Barras E, Reith W, Santos MA, and Pierre P. MicroRNA-155 modulates the interleukin-1 signaling pathway in activated human monocyte-derived dendritic cells. *Proc Natl Acad Sci USA* 106: 2735-2740, 2009.
39. Chan MH, Carey AL, Watt MJ, and Febbraio MA. Cytokine gene expression in human skeletal muscle during concentric contraction: evidence that IL-8, like IL-6, is influenced by glycogen availability. *Am J Physiol Regul Integr Comp Physiol* 287: R322-R327, 2004.
40. Chazaud B, Sonnet C, Lafuste P, Bassez G, Rimaniol AC, Poron F, Authier FJ, Dreyfus PA, and Gherardi RK. Satellite cells attract monocytes and use macrophages as a support to escape apoptosis and enhance muscle growth. *J Cell Biol* 163: 1133-1143, 2003.
41. Chen CW, Montelatici E, Crisan M, Corselli M, Huard J, Lazhari L, and Peault B. Perivascular multi-lineage progenitor cells in human organs: regenerative units, cytokine sources or both? *Cytokine Growth Factor Rev* 20: 429-434, 2009.
42. Chen JF, Mandel EM, Thomson JM, Wu Q, Callis TE, Hammond SM, Conlon FL, and Wang DZ. The role of microRNA-1 and microRNA-133 in skeletal muscle proliferation and differentiation. *Nat Genet* 38: 228-233, 2006.
43. Chen SE, Jin B, and Li YP. TNF- α regulates myogenesis and muscle regeneration by activating p38 MAPK. *Am J Physiol Cell Physiol* 292: C1660-1671, 2007.
44. Chevrel G, Granet C, and Miossec P. Contribution of tumor necrosis factor α and interleukin (IL) 1 β to IL6 production, NF- κ B nuclear translocation, and class I MHC expression in muscle cells: in vitro regulation with specific cytokine inhibitors. *Ann Rheum Dis* 64: 1257-1262, 2005.
45. Christiansen T, Bruun JM, Paulsen SK, Olholm J, Overgaard K, Pedersen SB, and Richelsen B. Acute exercise increases circulating inflammatory markers in overweight and obese compared with lean subjects. *Eur J Appl Physiol* 113: 1635-1642, 2013.
46. De Bleecker JL, De Paep B, Vanwallegem IE, and Schroder JM. Differential expression of chemokines in inflammatory myopathies. *Neurology* 58: 1779-1785, 2002.
47. De Rossi M, Bernasconi P, Baggi F, de Waal Malefyt R, and Mantegazza R. Cytokines and chemokines are both expressed by human myoblasts: possible relevance for the immune pathogenesis of muscle inflammation. *Int Immunol* 12: 1329-1335, 2000.
48. de Waal Malefyt R, Abrams J, Bennett B, Figdor CG, and de Vries JE. Interleukin 10 (IL-10) inhibits cytokine synthesis by human monocytes: an autoregulatory role of IL-10 produced by monocytes. *J Exp Med* 174: 1209-1220, 1991.
49. Dean JL, Wait R, Mahtani KR, Sully G, Clark AR, and Saklatvala J. The 3' untranslated region of tumor necrosis factor α mRNA is a target of the mRNA-stabilizing factor HuR. *Mol Cell Biol* 21: 721-730, 2001.
50. Della Gatta PA, Cameron-Smith D, and Peake JM. Acute resistance exercise increases the expression of chemotactic factors within skeletal muscle. *Eur J Appl Physiol* 11: 2157-2167, 2014.
51. Della Gatta PA, Garnham AP, Peake JM, and Cameron-Smith D. Effect of exercise training on skeletal muscle cytokine expression in the elderly. *Brain Behav Immun* 39: 80-86, 2014.
52. Dennis RA, Trappe TA, Simpson P, Carroll C, Huang BE, Nagarajan R, Bearden E, Gurley C, Duff GW, Evans WJ, Kornman K, and Peterson CA. Interleukin-1 polymorphisms are associated with the inflammatory response in human muscle to acute resistance exercise. *J Physiol* 560: 617-626, 2004.
53. Dentelli P, Rosso A, Orso F, Olgasi C, Taverna D, and Brizzi MF. microRNA-222 controls neovascularization by regulating signal transducer and activator of transcription 5A expression. *Arterioscler Thromb Vasc Biol* 30: 1562-1568, 2010.
54. Dinarello CA, Ikejima T, Warner SJ, Orencole SF, Lonnemann G, Cannon JG, and Libby P. Interleukin 1 induces interleukin 1. I. Induction of circulating interleukin 1 in rabbits in vivo and in human mononuclear cells in vitro. *J Immunol* 139: 1902-1910, 1987.
55. Dittrich E, Rose-John S, Gerhartz C, Mullberg J, Stoyan T, Yasukawa K, Heinrich PC, and Graeve L. Identification of a region within the cytoplasmic domain of the interleukin-6 (IL-6) signal transducer gp130 important for ligand-induced endocytosis of the IL-6 receptor. *J Biol Chem* 269: 19014-19020, 1994.
56. Drenth J, Van Uum S, Van Deuren M, GJ P, Van Der Ven-Jongekrijg J, and Van Der Meer J. Endurance run increases circulating IL-6 and IL-1ra but downregulates ex vivo TNF- α and IL-1 β production. *J Appl Physiol* 79: 1497-1503, 1995.
57. Drenth JP, Krebbers RJ, Bijzet J, and van der Meer JW. Increased circulating cytokine receptors and ex vivo interleukin-1 receptor antagonist and interleukin-1 β production but decreased tumor necrosis factor- α production after a 5-km run. *Eur J Clin Invest* 28: 866-872, 1998.
58. Dubravec DB, Spriggs DR, Mannick JA, and Rodrick ML. Circulating human peripheral blood granulocytes synthesize and secrete tumor necrosis factor α . *Proc Natl Acad Sci USA* 87: 6758-6761, 1990.
59. El Gazzar M, and McCall CE. MicroRNAs distinguish translational from transcriptional silencing during endotoxin tolerance. *J Biol Chem* 285: 20940-20951, 2010.
60. Emanuelli B, Peraldi P, Filloux C, Chavey C, Freidinger K, Hilton DJ, Hotamisligil GS, and Van Obberghen E. SOCS-3 inhibits insulin signaling and is up-regulated in response to tumor necrosis factor- α in the adipose tissue of obese mice. *J Biol Chem* 276: 47944-47949, 2001.
61. Ensor JE, Wiener SM, McCrea KA, Viscardi RM, Crawford EK, and Hasday JD. Differential effects of hyperthermia on macrophage interleukin-6 and tumor necrosis factor- α expression. *Am J Physiol Cell Physiol* 266: C967-974, 1994.
62. Farmawati A, Kitajima Y, Nedachi T, Sato M, Kanzaki M, and Nagatomi R. Characterization of contraction-induced IL-6 up-regulation using contractile C2C12 myotubes. *Endocr J* 60: 137-147, 2013.

63. Febbraio MA, Steensberg A, Fischer CP, Keller C, Hiscock N, and Pedersen BK. IL-6 activates HSP72 gene expression in human skeletal muscle. *Biochem Biophys Res Commun* 296: 1264-1266, 2002.
64. Febbraio MA, Steensberg A, Keller C, Starkie RL, Nielsen HB, Krstrup P, Ott P, Secher NH, and Pedersen BK. Glucose ingestion attenuates interleukin-6 release from contracting skeletal muscle in humans. *J Physiol* 549: 607-612, 2003.
65. Febbraio MA, Steensberg A, Starkie RL, McConell GK, and Kingwell BA. Skeletal muscle interleukin-6 and tumor necrosis factor- α release in healthy subjects and patients with type 2 diabetes at rest and during exercise. *Metabolism* 52: 939-944, 2003.
66. Fielding R, Manfredi T, Ding W, Fiatarone M, Evans W, and Cannon J. Acute phase response in exercise III. Neutrophil and IL-1 β accumulation in skeletal muscle. *Am J Physiol Regul Integr Comp Physiol* 265: R166-R172, 1993.
67. Fiorentino DF, Zlotnik A, Mosmann TR, Howard M, and O'Garra A. IL-10 inhibits cytokine production by activated macrophages. *J Immunol* 147: 3815-3822, 1991.
68. Fischer C, Hiscock N, Penkowa M, Basu S, Vessby B, Kallner A, Sjoberg L-B, and Pedersen B. Vitamin C and E supplementation inhibits the release of interleukin-6 from contracting human skeletal muscle. *J Physiol* 558: 633-645, 2004.
69. Frost R, Nystrom G, and Lang C. Multiple Toll-like receptor ligands induce an IL-6 transcriptional response in skeletal myocytes. *Am J Physiol Regul Integr Comp Physiol* 290: R773-784, 2006.
70. Frost RA, Nystrom GJ, and Lang CH. Epinephrine stimulates IL-6 expression in skeletal muscle and C2C12 myoblasts: role of c-Jun NH2-terminal kinase and histone deacetylase activity. *Am J Physiol Endocrinol Metab* 286: E809-817, 2004.
71. Frost RA, Nystrom GJ, and Lang CH. Lipopolysaccharide and proinflammatory cytokines stimulate interleukin-6 expression in C2C12 myoblasts: role of the Jun NH2-terminal kinase. *Am J Physiol Regul Integr Comp Physiol* 285: R1153-1164, 2003.
72. Frost RA, Nystrom GJ, and Lang CH. Lipopolysaccharide regulates proinflammatory cytokine expression in mouse myoblasts and skeletal muscle. *Am J Physiol Regul Integr Comp Physiol* 283: R698-709, 2002.
73. Gallucci S, Provenzano C, Mazzarelli P, Scuderi F, and Bartocioni E. Myoblasts produce IL-6 in response to inflammatory stimuli. *Int Immunol* 10: 267-273, 1998.
74. Ge Y, Waldemer RJ, Nalluri R, Nuzzi PD, and Chen J. RNAi screen reveals potentially novel roles of cytokines in myoblast differentiation. *PLoS One* 8: e68068, 2013.
75. Geyer BC, Ben Ari S, Barbash S, Kilbourne J, Mor TS, and Soreq H. Nicotinic stimulation induces Tristetraprolin overproduction and attenuates inflammation in muscle. *Biochim Biophys Acta* 1823: 368-378, 2012.
76. Gilchrist M, Thorsson V, Li B, Rust AG, Korb M, Roach JC, Kennedy K, Hai T, Bolouri H, and Aderem A. Systems biology approaches identify ATF3 as a negative regulator of Toll-like receptor 4. *Nature* 441: 173-178, 2006.
77. Giresi PG, Stevenson EJ, Theilhaber J, Koncarevic A, Parkington J, Fielding RA, and Kandarian SC. Identification of a molecular signature of sarcopenia. *Physiol Genomics* 21: 253-263, 2005.
78. Grabstein KH, Eisenman J, Shanebeck K, Rauch C, Srinivasan S, Fung V, Beers C, Richardson J, Schoenborn MA, and Ahdieh M. Cloning of a T cell growth factor that interacts with the β chain of the interleukin-2 receptor. *Science* 264: 965-968, 1994.
79. Graeve L, Korolenko TA, Hemmann U, Weiergraber O, Ditrlich E, and Heinrich PC. A complex of the soluble interleukin-6 receptor and interleukin-6 is internalized via the signal transducer gp130. *FEBS Lett* 399: 131-134, 1996.
80. Griffin CA, Apponi LH, Long KK, and Pavlath GK. Chemokine expression and control of muscle cell migration during myogenesis. *J Cell Sci* 123: 3052-3060, 2010.
81. Haahr PM, Pedersen BK, Fomsgaard A, Tvede N, Diamant M, Klarlund K, Halkjaer-Kristensen J, and Bendtzen K. Effect of physical exercise on in vitro production of interleukin 1, interleukin 6, tumour necrosis factor- α , interleukin 2 and interferon- γ . *Int J Sports Med* 12: 223-227, 1991.
82. Hamada K, Vannier E, Sackeck JM, Witsell AL, and Roubenoff R. Senescence of human skeletal muscle impairs the local inflammatory cytokine response to acute eccentric exercise. *FASEB J* 19: 264-266, 2005.
83. Haugen F, Norheim F, Lian H, Wensaas AJ, Dueland S, Berg O, Funderud A, Skalhogg BS, Raastad T, and Drevon CA. IL-7 is expressed and secreted by human skeletal muscle cells. *Am J Physiol Cell Physiol* 298: C807-816, 2010.
84. Heaney ML, and Golde DW. Soluble cytokine receptors. *Blood* 87: 847-857, 1996.
85. Henningsen J, Pedersen BK, and Kratchmarova I. Quantitative analysis of the secretion of the MCP family of chemokines by muscle cells. *Mol Biosyst* 7: 311-321, 2011.
86. Henningsen J, Rigbolt KT, Blagoev B, Pedersen BK, and Kratchmarova I. Dynamics of the skeletal muscle secretome during myoblast differentiation. *Mol Cell Proteomics* 2010.
87. Henson DA, Nieman DC, Blodgett AD, Butterworth DE, Utter A, Davis JM, Sonnenfeld G, Morton DS, Fagoaga OR, and Nehlsen-Cannarella SL. Influence of exercise mode and carbohydrate on the immune response to prolonged exercise. *Int J Sport Nutr* 9: 213-228, 1999.
88. Higuchi M, and Aggarwal BB. TNF induces internalization of the p60 receptor and shedding of the p80 receptor. *J Immunol* 152: 3550-3558, 1994.
89. Hiscock N, Chan MH, Bisucci T, Darby IA, and Febbraio MA. Skeletal myocytes are a source of interleukin-6 mRNA expression and protein release during contraction: evidence of fiber type specificity. *FASEB J* 18: 992-994, 2004.
90. Hoier B, Prats C, Qvortrup K, Pilegaard H, Bangsbo J, and Hellsten Y. Subcellular localization and mechanism of secretion of vascular endothelial growth factor in human skeletal muscle. *FASEB J* 27: 3496-3504, 2013.
91. Hoier B, Walker M, Passos M, Walker PJ, Green A, Bangsbo J, Askew CD, and Hellsten Y. Angiogenic response to passive movement and active exercise in individuals with peripheral arterial disease. *J Appl Physiol* 115: 1777-1787, 2013.
92. Holmes AG, Watt MJ, Carey AL, and Febbraio MA. Ionomycin, but not physiologic doses of epinephrine, stimulates skeletal muscle interleukin-6 mRNA expression and protein release. *Metabolism* 53: 1492-1495, 2004.
93. Horii Y, Muraguchi A, Suematsu S, Matsuda T, Yoshizaki K, Hirano T, and Kishimoto T. Regulation of BSF-2/IL-6 production by human mononuclear cells. Macrophage-dependent synthesis of BSF-2/IL-6 by T cells. *J Immunol* 141: 1529-1535, 1988.
94. Hubal MJ, Chen TC, Thompson PD, and Clarkson PM. Inflammatory gene changes associated with the repeated-bout effect. *Am J Physiol Regul Integr Comp Physiol* 294: R1628-1637, 2008.

95. Huey KA, and Meador BM. Contribution of IL-6 to the Hsp72, Hsp25, and α B-crystallin responses to inflammation and exercise training in mouse skeletal and cardiac muscle. *J Appl Physiol* 105: 1830-1836, 2008.
96. Jacquemin V, Butler-Browne GS, Furling D, and Mouly V. IL-13 mediates the recruitment of reserve cells for fusion during IGF-1-induced hypertrophy of human myotubes. *J Cell Sci* 120: 670-681, 2007.
97. Jiang LQ, Franck N, Egan B, Sjogren RJ, Katayama M, Duque-Guimaraes D, Arner P, Zierath JR, and Krook A. Autocrine role of interleukin-13 on skeletal muscle glucose metabolism in type 2 diabetic patients involves microRNA let-7. *Am J Physiol Endocrinol Metab* 305: E1359-1366, 2013.
98. Jing Q, Huang S, Guth S, Zarubin T, Motoyama A, Chen J, Di Padova F, Lin SC, Gram H, and Han J. Involvement of microRNA in AU-rich element-mediated mRNA instability. *Cell* 120: 623-634, 2005.
99. Kakani MW, Peake J, Brenu EW, Simmonds M, Gray B, and Marshall-Gradisnik SM. T helper cell cytokine profiles after endurance exercise. *J Interferon Cytokine Res* 34: 699-706, 2014.
100. Kaplanski G, Farnarier C, Kaplanski S, Porat R, Shapiro L, Bongrand P, and Dinarello CA. Interleukin-1 induces interleukin-8 secretion from endothelial cells by a juxtacrine mechanism. *Blood* 84: 4242-4248, 1994.
101. Katsanou V, Papadaki O, Milatos S, Blackshear PJ, Anderson P, Kollias G, and Kontoyiannis DL. HuR as a negative post-transcriptional modulator in inflammation. *Mol Cell* 19: 777-789, 2005.
102. Keller C, Hellsten Y, Steensberg A, and Pedersen BK. Differential regulation of IL-6 and TNF- α via calcineurin in human skeletal muscle cells. *Cytokine* 36: 141-147, 2006.
103. Keller C, Keller P, Marshal S, and Pedersen BK. IL-6 gene expression in human adipose tissue in response to exercise - effect of carbohydrate ingestion. *J Physiol* 550: 927-931, 2003.
104. Keller P, Penkowa M, Keller C, Steensberg A, Fischer CP, Giralt M, Hidalgo J, and Pedersen BK. Interleukin-6 receptor expression in contracting human skeletal muscle: regulating role of IL-6. *FASEB J* 19: 1181-1183, 2005.
105. Kim HK, Lee YS, Sivaprasad U, Malhotra A, and Dutta A. Muscle-specific microRNA miR-206 promotes muscle differentiation. *J Cell Biol* 174: 677-687, 2006.
106. Kim JA, Park HS, Kang SR, Park KI, Lee DH, Nagappan A, Shin SC, Lee WS, Kim EH, and Kim GS. Suppressive effect of flavonoids from Korean citrus aurantium L. on the expression of inflammatory mediators in L6 skeletal muscle cells. *Phytother Res* 26: 1906-12, 2012.
107. Kosmidou I, Vassilakopoulos T, Xagorari A, Zakyntinos S, Papapetropoulos A, and Roussos C. Production of interleukin-6 by skeletal myotubes: role of reactive oxygen species. *Am J Respir Cell Mol Biol* 26: 587-593, 2002.
108. Kovac A, Erickson MA, and Banks WA. Brain microvascular pericytes are immunoactive in culture: cytokine, chemokine, nitric oxide, and LRP-1 expression in response to lipopolysaccharide. *J Neuroinflammation* 8: 139-148, 2011.
109. Krogh-Madsen R, Plomgaard P, Moller K, Mittendorfer B, and Pedersen BK. Influence of TNF- α and IL-6 infusions on insulin sensitivity and expression of IL-18 in humans. *Am J Physiol Endocrinol Metab* 291: E108-114, 2006.
110. Kurek JB, Bower JJ, Romanella M, Koentgen F, Murphy M, and Austin L. The role of leukemia inhibitory factor in skeletal muscle regeneration. *Muscle Nerve* 20: 815-822, 1997.
111. Kurt-Jones EA, Fiers W, and Pober JS. Membrane interleukin 1 induction on human endothelial cells and dermal fibroblasts. *J Immunol* 139: 2317-2324, 1987.
112. Kvernmo H, Olsen JO, and Osterud B. Changes in blood cell response following strenuous physical exercise. *Eur J Appl Physiol Occup Physiol* 64: 318-322, 1992.
113. Lancaster GL, Halson SL, Khan Q, Drysdale P, Wallace F, Jeukendrup AE, Drayson MT, and Gleeson M. Effects of acute exhaustive exercise and chronic exercise training on type 1 and type 2 T lymphocytes. *Exerc Immunol Rev* 10: 91-106, 2004.
114. Langberg H, Olesen JL, Gemmer C, and Kjaer M. Substantial elevation of interleukin-6 concentration in peritendinous tissue, in contrast to muscle, following prolonged exercise in humans. *J Physiol* 542: 985-990, 2002.
115. Langen RC, Schols AM, Kelders MC, Wouters EF, and Janssen-Heininger YM. Inflammatory cytokines inhibit myogenic differentiation through activation of nuclear factor- κ B. *FASEB J* 15: 1169-1180, 2001.
116. Lauritzen HP, Brandauer J, Schjerling P, Koh HJ, Treebak JT, Hirshman MF, Galbo H, and Goodyear LJ. Contraction and AICAR stimulate IL-6 vesicle depletion from skeletal muscle fibers in vivo. *Diabetes* 62: 3081-3092, 2013.
117. LaVoy EC, Bosch JA, Lowder TW, and Simpson RJ. Acute aerobic exercise in humans increases cytokine expression in CD27(-) but not CD27(+) CD8(+) T-cells. *Brain Behav Immun* 27: 54-62, 2013.
118. Le JM, Weinstein D, Gubler U, and Vilcek J. Induction of membrane-associated interleukin 1 by tumor necrosis factor in human fibroblasts. *J Immunol* 138: 2137-2142, 1987.
119. Lee WJ. IGF-I exerts an anti-inflammatory effect on skeletal muscle cells through down-regulation of TLR4 signaling. *Immune Netw* 11: 223-226, 2011.
120. Leggate M, Nowell MA, Jones SA, and Nimmo MA. The response of interleukin-6 and soluble interleukin-6 receptor isoforms following intermittent high intensity and continuous moderate intensity cycling. *Cell Stress Chaperones* 15: 827-833, 2010.
121. Li Y, and Huard J. Differentiation of muscle-derived cells into myofibroblasts in injured skeletal muscle. *Am J Pathol* 161: 895-907, 2002.
122. Li YP. TNF- α is a mitogen in skeletal muscle. *Am J Physiol Cell Physiol* 285: C370-376, 2003.
123. Liebler JM, Kunkel SL, Burdick MD, Standiford TJ, Rolfe MW, and Strieter RM. Production of IL-8 and monocyte chemoattractant peptide-1 by peripheral blood monocytes. Disparate responses to phytohemagglutinin and lipopolysaccharide. *J Immunol* 152: 241-249, 1994.
124. Louis E, Raue U, Yang Y, Jemiolo B, and Trappe S. Time course of proteolytic, cytokine, and myostatin gene expression after acute exercise in human skeletal muscle. *J Appl Physiol* 103: 1744-1751, 2007.
125. Luo G, Hershko DD, Robb BW, Wray CJ, and Hasselgren PO. IL-1 β stimulates IL-6 production in cultured skeletal muscle cells through activation of MAP kinase signaling pathway and NF- κ B. *Am J Physiol Regul Integr Comp Physiol* 284: R1249-1254, 2003.
126. Lyngso D, Simonsen L, and Bulow J. Interleukin-6 production in human subcutaneous abdominal adipose tissue: the effect of exercise. *J Physiol* 543: 373-378, 2002.

127. Ma F, Liu X, Li D, Wang P, Li N, Lu L, and Cao X. MicroRNA-4661 upregulates IL-10 expression in TLR-triggered macrophages by antagonizing RNA-binding protein tristetraprolin-mediated IL-10 mRNA degradation. *J Immunol* 184: 6053-6059, 2010.
128. Makris AC, Sotzios Y, Zhou Z, Makropoulou M, Papapetropoulos N, Zacharatos P, Pyriochou A, Roussos C, Papapetropoulos A, and Vassilakopoulos T. Nitric oxide stimulates interleukin-6 production in skeletal myotubes. *J Interferon Cytokine Res* 30: 321-327, 2010.
129. Malm C, Nyberg P, Engstrom M, Sjodin B, Lenkei R, Ekblom B, and Lundberg I. Immunological changes in human skeletal muscle and blood after eccentric exercise and multiple biopsies. *J Physiol* 529: 243-262, 2000.
130. Malm C, Sjodin TL, Sjoberg B, Lenkei R, Renstrom P, Lundberg IE, and Ekblom B. Leukocytes, cytokines, growth factors and hormones in human skeletal muscle and blood after uphill or downhill running. *J Physiol* 556: 983-1000, 2004.
131. Mannel DN, Moore RN, and Mergenhagen SE. Macrophages as a source of tumoricidal activity (tumor-necrotizing factor). *Infect Immun* 30: 523-530, 1980.
132. Marie C, Pitton C, Fitting C, and Cavaillon JM. IL-10 and IL-4 synergize with TNF- α to induce IL-1ra production by human neutrophils. *Cytokine* 8: 147-151, 1996.
133. Marino M, Scuderi F, Provenzano C, Scheller J, Rose-John S, and Bartocioni E. IL-6 regulates MCP-1, ICAM-1 and IL-6 expression in human myoblasts. *J Neuroimmunol* 196: 41-48, 2008.
134. Marker T, Sell H, Zillessen P, Glode A, Kriebel J, Ouwens DM, Pattyn P, Ruige J, Famulla S, Roden M, Eckel J, and Habich C. Heat shock protein 60 as a mediator of adipose tissue inflammation and insulin resistance. *Diabetes* 61: 615-625, 2012.
135. Markovitch D, Tyrrell RM, and Thompson D. Acute moderate-intensity exercise in middle-aged men has neither an anti- nor proinflammatory effect. *J Appl Physiol* 105: 260-265, 2008.
136. Martel-Pelletier J, McCollum R, and Pelletier JP. The synthesis of IL-1 receptor antagonist (IL-1ra) by synovial fibroblasts is markedly increased by the cytokines TNF- α and IL-1. *Biochim Biophys Acta* 1175: 302-305, 1993.
137. Matthews VB, Astrom MB, Chan MH, Bruce CR, Krabbe KS, Prelovsek O, Akerstrom T, Yfanti C, Broholm C, Mortensen OH, Penkowa M, Hojman P, Zankari A, Watt MJ, Bruunsgaard H, Pedersen BK, and Febbraio MA. Brain-derived neurotrophic factor is produced by skeletal muscle cells in response to contraction and enhances fat oxidation via activation of AMP-activated protein kinase. *Diabetologia* 52: 1409-1418, 2009.
138. McKay BR, De Lisio M, Johnston AP, O'Reilly CE, Phillips SM, Tarnopolsky MA, and Parise G. Association of interleukin-6 signalling with the muscle stem cell response following muscle-lengthening contractions in humans. *PLoS One* 4: e6027, 2009.
139. McKinsey TA, Zhang CL, Lu J, and Olson EN. Signal-dependent nuclear export of a histone deacetylase regulates muscle differentiation. *Nature* 408: 106-111, 2000.
140. McMullen MR, Cocuzzi E, Hatzoglou M, and Nagy LE. Chronic ethanol exposure increases the binding of HuR to the TNF α 3'-untranslated region in macrophages. *J Biol Chem* 278: 38333-38341, 2003.
141. Melo F, Carey DJ, and Brandan E. Extracellular matrix is required for skeletal muscle differentiation but not myogenin expression. *J Cell Biochem* 62: 227-239, 1996.
142. Meng X, Banerjee A, Ao L, Meldrum DR, Cain BS, Shames BD, and Harken AH. Inhibition of myocardial TNF- α production by heat shock. A potential mechanism of stress-induced cardioprotection against postischemic dysfunction. *Ann NY Acad Sci* 874: 69-82, 1999.
143. Miura P, Amirouche A, Clow C, Belanger G, and Jasmin BJ. Brain-derived neurotrophic factor expression is repressed during myogenic differentiation by miR-206. *J Neurochem* 120: 230-238, 2012.
144. Nagaraju K, Raben N, Merritt G, Loeffler L, Kirk K, and Plotz P. A variety of cytokines and immunologically relevant surface molecules are expressed by normal human skeletal muscle cells under proinflammatory stimuli. *Clin Exp Immunol* 113: 407-414, 1998.
145. Naka T, Narazaki M, and Kishimoto T. Negative-feedback regulation of cytokine signals. In: *Cytokine inhibitors*, edited by Ciliberto G. New York: Marcel Dekker, 2000, p. 241-260.
146. Nedachi T, Fujita H, and Kanzaki M. Contractile C2C12 myotube model for studying exercise-inducible responses in skeletal muscle. *Am J Physiol Endocrinol Metab* 295: E1191-E1204, 2008.
147. Neubauer O, Sabapathy S, Ashton KJ, Desbrow B, Peake JM, Lazarus R, Wessner B, Cameron-Smith D, Wagner KH, Haseler LJ, and Bulmer AC. Time course-dependent changes in the transcriptome of human skeletal muscle during recovery from endurance exercise: from inflammation to adaptive remodeling. *J Appl Physiol* 116: 274-287, 2014.
148. Nguyen HX, and Tidball JG. Null mutation of gp91phox reduces muscle membrane lysis during muscle inflammation in mice. *J Physiol* 553: 833-841, 2003.
149. Nielsen AR, Mounier R, Plomgaard P, Mortensen OH, Penkowa M, Speersneider T, Pilegaard H, and Pedersen BK. Expression of interleukin-15 in human skeletal muscle effect of exercise and muscle fibre type composition. *J Physiol* 584: 305-312, 2007.
150. Nieman DC, Davis JM, Brown VA, Henson DA, Dumke CL, Utter AC, Vinci DM, Downs MF, Smith JC, Carson J, Brown A, McAnulty SR, and McAnulty LS. Influence of carbohydrate ingestion on immune changes after 2 h of intensive resistance training. *J Appl Physiol* 96: 1292-1298, 2004.
151. Nieman DC, Davis JM, Henson DA, Gross SJ, Dumke CL, Utter AC, Vinci DM, Carson JA, Brown A, McAnulty SR, McAnulty LS, and Triplett NT. Muscle cytokine mRNA changes after 2.5 h of cycling: influence of carbohydrate. *Med Sci Sports Exerc* 37: 1283-1290, 2005.
152. Nieman DC, Davis JM, Henson DA, Walberg-Rankin J, Shute M, Dumke CL, Utter AC, Vinci DM, Carson JA, Brown A, Lee WJ, McAnulty SR, and McAnulty LS. Carbohydrate ingestion influences skeletal muscle cytokine mRNA and plasma cytokine levels after a 3-h run. *J Appl Physiol* 94: 1917-1925, 2003.
153. Nieman DC, Dumke CL, Henson DA, McAnulty SR, Gross SJ, and Lind RH. Muscle damage is linked to cytokine changes following a 160-km race. *Brain Behav Immun* 19: 398-403, 2005.
154. Nieman DC, Henson DA, Davis JM, Angela Murphy E, Jenkins DP, Gross SJ, Carmichael MD, Quindry JC, Dumke CL, Utter AC, McAnulty SR, McAnulty LS, Triplett NT, and Mayer EP. Quercetin's influence on exercise-induced changes in plasma cytokines and muscle and leukocyte cytokine mRNA. *J Appl Physiol* 103: 1728-1735, 2007.

155. Nieman DC, Henson DA, Smith LL, Utter AC, Vinci DM, Davis JM, Kaminsky DE, and Shute M. Cytokine changes after a marathon race. *J Appl Physiol* 91: 109-114, 2001.
156. Nieman DC, Konrad M, Henson DA, Kennerly K, Shanely RA, and Wallner-Liebmann SJ. Variance in the acute inflammatory response to prolonged cycling is linked to exercise intensity. *J Interferon Cytokine Res* 32: 12-17, 2012.
157. Nieman DC, Luo B, Dreau D, Henson DA, Shanely RA, Dew D, and Meaney MP. Immune and inflammation responses to a 3-day period of intensified running versus cycling. *Brain Behav Immun* 39: 180-185, 2014.
158. Nybo L, Nielsen B, Pedersen BK, Moller K, and Secher NH. Interleukin-6 release from the human brain during prolonged exercise. *J Physiol* 542: 991-995, 2002.
159. Ohno Y, Yamada S, Sugiura T, Ohira Y, Yoshioka T, and Goto K. Possible role of NF- κ B signals in heat stress-associated increase in protein content of cultured C2C12 cells. *Cells Tissues Organs* 194: 363-370, 2011.
160. Ostrowski K, Rohde T, Asp S, Schjerling P, and Pedersen BK. Pro- and anti-inflammatory cytokine balance in strenuous exercise in humans. *J Physiol* 515: 287-291, 1999.
161. Ostrowski K, Rohde T, Zacho M, Asp S, and Pedersen B. Evidence that interleukin-6 is produced in human skeletal muscle during prolonged running. *J Physiol* 508: 949-953, 1998.
162. Otis JS, Niccoli S, Hawdon N, Sarvas JL, Frye MA, Chicco AJ, and Lees SJ. Pro-inflammatory mediation of myoblast proliferation. *PLoS One* 9: e92363, 2014.
163. Pang G, Couch L, Batey R, Clancy R, and Cripps A. GM-CSF, IL-1 α , IL-1 β , IL-6, IL-8, IL-10, ICAM-1 and VCAM-1 gene expression and cytokine production in human duodenal fibroblasts stimulated with lipopolysaccharide, IL-1 α and TNF- α . *Clin Exp Immunol* 96: 437-443, 1994.
164. Peake JM, Suzuki K, Hordern M, Wilson G, Nosaka K, and Coombes JS. Plasma cytokine changes in relation to exercise intensity and muscle damage. *Eur J Appl Physiol* 95: 514-521, 2005.
165. Peake JM, Wilson G, Hordern M, Suzuki K, Nosaka K, Yamaya K, Mackinnon L, and Coombes J. Changes in neutrophil receptor expression, degranulation and respiratory burst activity after moderate and high intensity exercise. *J Appl Physiol* 97: 612-618, 2004.
166. Pedersen BK. Muscles and their myokines. *J Exp Biol* 214: 337-346, 2011.
167. Pedersen BK, and Febbraio MA. Muscle as an endocrine organ: focus on muscle-derived interleukin-6. *Physiol Rev* 88: 1379-1406, 2008.
168. Pedersen BK, and Febbraio MA. Muscles, exercise and obesity: skeletal muscle as a secretory organ. *Nat Rev Endocrinol* 8: 457-465, 2012.
169. Perdiguero E, Ruiz-Bonilla V, Serrano AL, and Munoz-Canoves P. Genetic deficiency of p38 α reveals its critical role in myoblast cell cycle exit: the p38 α -JNK connection. *Cell Cycle* 6: 1298-1303, 2007.
170. Perry MM, Moschos SA, Williams AE, Shepherd NJ, Larner-Svensson HM, and Lindsay MA. Rapid changes in microRNA-146a expression negatively regulate the IL-1 β -induced inflammatory response in human lung alveolar epithelial cells. *J Immunol* 180: 5689-5698, 2008.
171. Petersen AM, and Pedersen BK. The anti-inflammatory effect of exercise. *J Appl Physiol* 98: 1154-1162, 2005.
172. Peterson J, and Pizza F. Cytokines derived from cultured skeletal muscle cells after mechanical strain promote neutrophil chemotaxis in vitro. *J Appl Physiol* 106: 130-137, 2009.
173. Phillips MD, Flynn MG, McFarlin BK, Stewart LK, Timmerman KL, and Ji H. Resistive exercise blunts LPS-stimulated TNF- α and IL-1 β . *Int J Sports Med* 29: 102-109, 2008.
174. Pizza FX, McLoughlin TJ, McGregor SJ, Calomeni EP, and Gunning WT. Neutrophils injure cultured skeletal myotubes. *Am J Physiol Cell Physiol* 281: C335-341, 2001.
175. Plomgaard P, Penkowa M, and Pedersen BK. Fiber type specific expression of TNF- α , IL-6 and IL-18 in human skeletal muscles. *Exerc Immunol Rev* 11: 53-63, 2005.
176. Ponte AL, Marais E, Gallay N, Langonne A, Delorme B, Herval O, Charbord P, and Domenech J. The in vitro migration capacity of human bone marrow mesenchymal stem cells: comparison of chemokine and growth factor chemotactic activities. *Stem Cells* 25: 1737-1745, 2007.
177. Porteu F, and Hieblot C. Tumor necrosis factor induces a selective shedding of its p75 receptor from human neutrophils. *J Biol Chem* 269: 2834-2840, 1994.
178. Prelovsek O, Mars T, Jevsek M, Podbregar M, and Grubic Z. High dexamethasone concentration prevents stimulatory effects of TNF- α and LPS on IL-6 secretion from the precursors of human muscle regeneration. *Am J Physiol Regul Integr Comp Physiol* 291: R1651-1656, 2006.
179. Pue CA, Mortensen RF, Marsh CB, Pope HA, and Wewers MD. Acute phase levels of C-reactive protein enhance IL-1 β and IL-1ra production by human blood monocytes but inhibit IL-1 β and IL-1ra production by alveolar macrophages. *J Immunol* 156: 1594-1600, 1996.
180. Quinn LS, Strait-Bodey L, Anderson BG, Argiles JM, and Havel PJ. Interleukin-15 stimulates adiponectin secretion by 3T3-L1 adipocytes: evidence for a skeletal muscle-to-fat signaling pathway. *Cell Biol Int* 29: 449-457, 2005.
181. Rajasingh J, Bord E, Luedemann C, Asai J, Hamada H, Thorne T, Qin G, Goukassian D, Zhu Y, Losordo DW, and Kishore R. IL-10-induced TNF- α mRNA destabilization is mediated via IL-10 suppression of p38 MAP kinase activation and inhibition of HuR expression. *FASEB J* 20: 2112-2114, 2006.
182. Raschke S, Eckardt K, Bjorklund Holven K, Jensen J, and Eckel J. Identification and validation of novel contraction-regulated myokines released from primary human skeletal muscle cells. *PLoS One* 8: e62008, 2013.
183. Raschke S, and Eckel J. Adipo-myokines: two sides of the same coin--mediators of inflammation and mediators of exercise. *Mediators Inflamm* 2013: 320724, 2013.
184. Reihmane D, Jurka A, Tretjakovs P, and Dela F. Increase in IL-6, TNF- α , and MMP-9, but not sICAM-1, concentrations depends on exercise duration. *Eur J Appl Physiol* 113: 851-858, 2013.
185. Rhind SG, Castellani JW, Brenner IK, Shephard RJ, Zamecnik J, Montain SJ, Young AJ, and Shek PN. Intracellular monocyte and serum cytokine expression is modulated by exhausting exercise and cold exposure. *Am J Physiol Regul Integr Comp Physiol* 281: R66-R75, 2001.
186. Riechman SE, Balasekaran G, Roth SM, and Ferrell RE. Association of interleukin-15 protein and interleukin-15 receptor genetic variation with resistance exercise training responses. *J Appl Physiol* 97: 2214-2219, 2004.

187. Riedemann NC, Guo RF, Hollmann TJ, Gao H, Neff TA, Reuben JS, Speyer CL, Sarma JV, Wetsel RA, Zetoune FS, and Ward PA. Regulatory role of C5a in LPS-induced IL-6 production by neutrophils during sepsis. *FASEB J* 18: 370-372, 2004.
188. Rippe C, Blimline M, Magerko KA, Lawson BR, LaRocca TJ, Donato AJ, and Seals DR. MicroRNA changes in human arterial endothelial cells with senescence: relation to apoptosis, eNOS and inflammation. *Exp Gerontol* 47: 45-51, 2012.
189. Rivier A, Pene J, Chanez P, Anshelme F, Caillaud C, Prefaut C, Godard P, and Bousquet J. Release of cytokines by blood monocytes during strenuous exercise. *International Journal of Sports Medicine* 15: 192-198, 1994.
190. Rojas Vega S, Knicker A, Hollmann W, Bloch W, and Struder HK. Effect of resistance exercise on serum levels of growth factors in humans. *Horm Metab Res* 42: 982-986, 2010.
191. Rosendal L, Sogaard K, Kjaer M, Sjogaard G, Langberg H, and Kristiansen J. Increase in interstitial interleukin-6 of human skeletal muscle with repetitive low-force exercise. *J Appl Physiol* 98: 477-481, 2005.
192. Ross ML, Halson SL, Suzuki K, Garnham A, Hawley JA, Cameron-Smith D, and Peake JM. Cytokine responses to carbohydrate ingestion during recovery from exercise-induced muscle injury. *J Interferon Cytokine Res* 30: 329-337, 2010.
193. Sachidanandan C, Sambasivan R, and Dhawan J. Tristetraprolin and LPS-inducible CXC chemokine are rapidly induced in presumptive satellite cells in response to skeletal muscle injury. *J Cell Sci* 115: 2701-2712, 2002.
194. Salomonsson S, and Lundberg IE. Cytokines in idiopathic inflammatory myopathies. *Autoimmunity* 39: 177-190, 2006.
195. Scheler M, Irmeler M, Lehr S, Hartwig S, Staiger H, Al-Hasani H, Beckers J, de Angelis MH, Haring HU, and Weigert C. Cytokine response of primary human myotubes in an in vitro exercise model. *Am J Physiol Cell Physiol* 305: C877-886, 2013.
196. Schindler R, Mancilla J, Endres S, Ghorbani R, Clark SC, and Dinarello CA. Correlations and interactions in the production of interleukin-6 (IL-6), IL-1, and tumor necrosis factor (TNF) in human blood mononuclear cells: IL-6 suppresses IL-1 and TNF. *Blood* 75: 40-47, 1990.
197. Serrano AL, Baeza-Raja B, Perdiguero E, Jardi M, and Munoz-Canoves P. Interleukin-6 is an essential regulator of satellite cell-mediated skeletal muscle hypertrophy. *Cell Metab* 7: 33-44, 2008.
198. Sharma A, Kumar M, Aich J, Hariharan M, Brahmachari SK, Agrawal A, and Ghosh B. Posttranscriptional regulation of interleukin-10 expression by hsa-miR-106a. *Proc Natl Acad Sci USA* 106: 5761-5766, 2009.
199. Shin JN, Kim I, Lee JS, Koh GY, Lee ZH, and Kim HH. A novel zinc finger protein that inhibits osteoclastogenesis and the function of tumor necrosis factor receptor-associated factor 6. *J Biol Chem* 277: 8346-8353, 2002.
200. Simbertsev A, and Kozlov I. Cytokine system. In: *Mechanical stretch and cytokines* [e-book], edited by Kamkin A, and Kiseleva I. Dordrecht: Springer Science+Business Media B.V., 2012, p. 1-6.
201. Smith CA, Stauber F, Waters C, Alway SE, and Stauber WT. Transforming growth factor- β following skeletal muscle strain injury in rats. *J Appl Physiol* 102: 755-761, 2007.
202. Smits HH, Grunberg K, Derijk RH, Sterk PJ, and Hiemstra PS. Cytokine release and its modulation by dexamethasone in whole blood following exercise. *Clin Exp Immunol* 111: 463-468, 1998.
203. Snell ES, and Atkins E. The presence of endogenous pyrogen in normal rabbit tissues. *J Exp Med* 121: 1019-1038, 1965.
204. Spangenburg EE, Brown DA, Johnson MS, and Moore RL. Exercise increases SOCS-3 expression in rat skeletal muscle: potential relationship to IL-6 expression. *J Physiol* 572: 839-848, 2006.
205. Stamou P, and Kontoyiannis DL. Posttranscriptional regulation of TNF mRNA: a paradigm of signal-dependent mRNA utilization and its relevance to pathology. *Curr Dir Autoimmun* 11: 61-79, 2010.
206. Starkie R, Ostrowski SR, Jauffred S, Febbraio M, and Pedersen BK. Exercise and IL-6 infusion inhibit endotoxin-induced TNF- α production in humans. *FASEB J* 17: 884-886, 2003.
207. Starkie RL, Angus DJ, Rolland J, Hargreaves M, and Febbraio MA. Effect of prolonged, submaximal exercise and carbohydrate ingestion on monocyte intracellular cytokine production in humans. *J Physiol* 528: 647-655, 2000.
208. Starkie RL, Arkinstall MJ, Koukoulas I, Hawley JA, and Febbraio MA. Carbohydrate ingestion attenuates the increase in plasma interleukin-6, but not skeletal muscle interleukin-6 mRNA, during exercise in humans. *J Physiol* 533: 585-591, 2001.
209. Starkie RL, Hargreaves M, Rolland J, and Febbraio M. Heat stress, cytokines and the immune response to exercise. *Brain Behav Immun* 19: 404-412, 2005.
210. Starkie RL, Rolland J, Angus DJ, Anderson MJ, and Febbraio MA. Circulating monocytes are not the source of elevations in plasma IL-6 and TNF- α levels after prolonged running. *Am J Physiol Cell Physiol* 280: C769-C774, 2001.
211. Steensberg A, Febbraio MA, Osada T, Schjerling P, van Hall G, Saltin B, and Pedersen BK. Interleukin-6 production in contracting human skeletal muscle is influenced by pre-exercise muscle glycogen content. *J Physiol* 537: 633-639, 2001.
212. Steensberg A, Keller C, Starkie R, Osada T, Febbraio M, and Pedersen B. IL-6 and TNF- α expression in, and release from, contracting human skeletal muscle. *Am J Physiol Endocrinol Metab* 283: E1272-E1278, 2002.
213. Steensberg A, Toft AD, Bruunsgaard H, Sandmand M, Halkjaer-Kristensen J, and Pedersen BK. Strenuous exercise decreases the percentage of type 1 T cells in the circulation. *J Appl Physiol* 91: 1708-1712, 2001.
214. Steensberg A, Toft AD, Schjerling P, Halkjaer-Kristensen J, and Pedersen BK. Plasma interleukin-6 during strenuous exercise: role of epinephrine. *Am J Physiol Cell Physiol* 281: C1001-C1004, 2001.
215. Steensberg A, van Hall G, Osada T, Sacchetti M, Saltin B, and Pedersen BK. Production of interleukin-6 in contracting human skeletal muscles can account for the exercise-induced increase in plasma interleukin-6. *J Physiol* 529: 237-242, 2000.
216. Stow JL, and Murray RZ. Intracellular trafficking and secretion of inflammatory cytokines. *Cytokine Growth Factor Rev* 24: 227-239, 2013.
217. Struyf S, Van Collie E, Paemen L, Put W, Lenaerts JP, Proost P, Opdenakker G, and Van Damme J. Synergistic induction of MCP-1 and -2 by IL-1 β and interferons in fibroblasts and epithelial cells. *J Leukoc Biol* 63: 364-372, 1998.
218. Summan M, Warren GL, Mercer RR, Chapman R, Hulderman T, Van Rooijen N, and Simeonova PP. Macrophages and skeletal muscle regeneration: a clodronate-containing liposome depletion study. *Am J Physiol Regul Integr Comp Physiol* 290: R1488-1495, 2006.

219. Suzuki K, Nakaji S, Yamada M, Liu Q, Kurakake S, Okamura N, Kumae T, Umeda T, and Sugawara K. Impact of a competitive marathon race on systemic cytokine and neutrophil responses. *Med Sci Sports Exerc* 35: 348-355, 2003.
220. Suzuki K, Nakaji S, Yamada M, Totsuka M, Sato K, and Sugawara K. Systemic inflammatory response to exhaustive exercise. Cytokine kinetics. *Exerc Immunol Rev* 8: 6-48, 2002.
221. Suzuki K, Nakajima H, Ikeda K, Maezawa Y, Suto A, Takatori H, Saito Y, and Iwamoto I. IL-4-Stat6 signaling induces tristetraprolin expression and inhibits TNF- α production in mast cells. *J Exp Med* 198: 1717-1727, 2003.
222. Taganov KD, Boldin MP, Chang KJ, and Baltimore D. NF- κ B-dependent induction of microRNA miR-146, an inhibitor targeted to signaling proteins of innate immune responses. *Proc Natl Acad Sci USA* 103: 12481-12486, 2006.
223. Tiku K, Tiku ML, and Skosey JL. Interleukin 1 production by human polymorphonuclear neutrophils. *J Immunol* 136: 3677-3685, 1986.
224. Tili E, Michaille JJ, Cimino A, Costinean S, Dumitru CD, Adair B, Fabbri M, Alder H, Liu CG, Calin GA, and Croce CM. Modulation of miR-155 and miR-125b levels following lipopolysaccharide/TNF- α stimulation and their possible roles in regulating the response to endotoxin shock. *J Immunol* 179: 5082-5089, 2007.
225. Toft AD, Jensen LB, Bruunsgaard H, Ibfelt T, Halkjaer-Kristensen J, Febbraio M, and Pedersen BK. Cytokine response to eccentric exercise in young and elderly humans. *Am J Physiol Cell Physiol* 283: C289-C295, 2002.
226. Tomiya A, Aizawa T, Nagatomi R, Sensui H, and Kokubun S. Myofibers express IL-6 after eccentric exercise. *Am J Sports Med* 32: 503-508, 2004.
227. Trendelenburg AU, Meyer A, Jacobi C, Feige JN, and Glass DJ. TAK-1/p38/NF κ B signaling inhibits myoblast differentiation by increasing levels of Activin A. *Skelet Muscle* 2: 3, 2012.
228. van de Vyver M, and Myburgh KH. Variable inflammation and intramuscular STAT3 phosphorylation and myeloperoxidase levels after downhill running. *Scand J Med Sci Sports* 24: e360-71, 2014.
229. Vasudevan S, Tong Y, and Steitz JA. Switching from repression to activation: microRNAs can up-regulate translation. *Science* 318: 1931-1934, 2007.
230. Vona-Davis L, Frankenberry K, Lebedeva K, and McFadden DW. LPS and TNF- α induce SOCS-3 mRNA expression in cardiac myoblasts. *Shock* 21: 38-38, 2004.
231. Weinstock C, Konig D, Harnischmacher R, Keul J, Berg A, and Northoff H. Effect of exhaustive exercise stress on the cytokine response. *Med Sci Sports Exerc* 29: 345-354, 1997.
232. Welc SS, and Clanton TL. The regulation of interleukin-6 implicates skeletal muscle as an integrative stress sensor and endocrine organ. *Exp Physiol* 98: 359-371, 2013.
233. Welc SS, Judge AR, and Clanton TL. Skeletal muscle interleukin-6 regulation in hyperthermia. *Am J Physiol Cell Physiol* 305: C406-413, 2013.
234. Welc SS, Phillips NA, Oca-Cossio J, Wallet SM, Chen DL, and Clanton TL. Hyperthermia increases interleukin-6 in mouse skeletal muscle. *Am J Physiol Cell Physiol* 303: C455-466, 2012.
235. Whitham M, Chan MH, Pal M, Matthews VB, Prelovsek O, Lunke S, El-Osta A, Broenneke H, Alber J, Bruning JC, Wunderlich FT, Lancaster GI, and Febbraio MA. Contraction-induced IL-6 gene transcription in skeletal muscle is regulated by c-jun terminal kinase/Activator protein -1. *J Biol Chem* 287: 10771-10779, 2012.
236. Xing Z, Gauldie J, Cox G, Baumann H, Jordana M, Lei XF, and Achong MK. IL-6 is an antiinflammatory cytokine required for controlling local or systemic acute inflammatory responses. *J Clin Invest* 101: 311-320, 1998.
237. Yahiaoui L, Gvozdic D, Daneliou G, Mack M, and Petrof BJ. CC chemokines directly regulate myoblast responses to skeletal muscle injury. *J Physiol* 586: 3991-4004, 2008.
238. Yan SF, Tritto I, Pinsky D, Liao H, Huang J, Fuller G, Brett J, May L, and Stern D. Induction of interleukin 6 (IL-6) by hypoxia in vascular cells. Central role of the binding site for nuclear factor-IL-6. *J Biol Chem* 270: 11463-11471, 1995.
239. Zaldivar F, Wang-Rodriguez J, Nemet D, Schwindt C, Galasetti P, Mills PJ, Wilson LD, and Cooper DM. Constitutive pro- and anti-inflammatory cytokine and growth factor response to exercise in leukocytes. *J Appl Physiol* 100: 1124-1133, 2006.
240. Zhang L, Lee JE, Wilusz J, and Wilusz CJ. The RNA-binding protein CUGBP1 regulates stability of tumor necrosis factor mRNA in muscle cells: implications for myotonic dystrophy. *J Biol Chem* 283: 22457-22463, 2008.