

Influence of age and physical fitness on miRNA-21, TGF- β and its receptors in leukocytes of healthy women

Barbara Halper¹, Marlene Hofmann², Stefan Oesen¹, Bernhard Franzke¹, Petra Stuparits³, Claudia Vidotto⁴, Harald Tschan², Norbert Bachl², Eva-Maria Strasser⁵, Michael Quittan⁵, Karl-Heinz Wagner^{1,6}, Barbara Wessner^{1,2}

¹ University of Vienna, Research Platform Active Ageing, Althanstraße 14, 1090 Vienna, Austria

² University of Vienna, Centre for Sport Science and University Sports, Department of Sport and Exercise Physiology, Auf der Schmelz 6, 1150 Vienna, Austria.

³ University of Vienna, Centre for Sport Science and University Sports, Department of Preventive and Rehabilitative Training Science, Auf der Schmelz 6, 1150 Vienna, Austria.

⁴ Study Lab G.m.b.H., Davidgasse 87-89, 1100 Vienna

⁵ Karl Landsteiner Institute for Remobilization and Functional Health/ Institute for Physical Medicine and Rehabilitation, Kaiser Franz Joseph Hospital, Social Medical Centre South, Kundratstrasse 3, 1100 Vienna, Austria.

⁶ University of Vienna, Faculty of Life Sciences, Department of Nutritional Science, Althanstraße 14, 1090 Vienna, Austria.

ABSTRACT

Rationale: The TGF- β superfamily has been shown to play an important role in a wide range of physiological as well as pathological processes including ageing, immune modulation, atherosclerosis and cancer development. The aim of the current study was to investigate (i) whether TGF- β signalling in peripheral blood mononuclear cells (PBMCs) would differ between young and old females and (ii) whether physical performance parameters of elderly women would be related to the expression of TGF- β or its receptors.

Methods: Sixteen healthy young (22-28 years; YF) and 90 healthy older (65-92 years; OF) females participated in the study. In addition to several components of health-related physical fitness, circulating CRP and TGF- β levels were determined together with the mRNA expression of TGF- β , TGF- β RI, TGF- β RII, and miRNA-21 (known to interfere with TGF- β signalling) in PBMCs.

Results: Physical fitness as determined by 6-minutes walking test (YF: median 932 (range 573-1254) m; OF: 360 (114-558) m), handgrip strength (YF: 32 (24-39) kg; OF: 18 (10-30) kg), relative isokinetic peak torque of knee extensors (YF: 1.9 (1.2-2.3) Nm/kg; OF: 1.0 (0.2-1.9) Nm/kg and flexors (YF: 1.1 (0.7-1.5) Nm/kg; OF: 0.5 (0.2-1.0) Nm/kg) was substantially lower in older women ($p < 0.001$ for all comparisons). These changes were paralleled by an increase in hs-CRP (YF: 0.9 (0.1-4.3) mg/L; OF: 2.3 (0.3-56.7) mg/L, $p < 0.001$). Serum levels of TGF- β and TGF- β mRNA levels from PBMCs did not differ between young and old women whereas, both TGF-

β RI/GAPDH (YF: 4.07 (1.38-14.60); OF: 2.08 (0.14-28.81); $p = 0.020$) and TGF- β RII/GAPDH levels (YF: 3.16 (1.14-10.25); OF: 1.71 (0.51-14.86); $p = 0.020$) were lower with respect to old age. In elderly women, only TGF- β RI expression correlated negatively with miRNA-21 expression in PBMCs ($\rho = -0.315$; $p = 0.004$). Interestingly, hs-CRP and miRNA correlated positively with handgrip strength ($\rho = 0.237$ and $\rho = 0.243$, $p < 0.05$), while none of the TGF- β -related parameters were related to physical performance.

Conclusion: The results suggest that age affects TGF- β signalling in leukocytes by altering the expression levels of its receptors. These changes seem to occur independently of physical fitness of old women.

Key Words: Inflamm-ageing, TGF- β Pathway, TGF- β receptors, microRNA-21, physical performance, Vienna Active Ageing Study

INTRODUCTION

Although the causes of human ageing are multifaceted, the molecular inflammation hypothesis of ageing implies that increased oxidative stress will lead to the activation of redox-sensitive transcription factors which in turn enhance the expression of pro-inflammatory genes in a variety of different cell types (8). As a consequence ageing is associated with a chronic inflammatory state, where pro-inflammatory factors such as tumour necrosis factor- α (TNF- α), interleukin-6 (IL-6) or C-reactive protein (CRP) are continuously present (over years) at levels higher than baseline but much lower than those found during acute inflammation (11, 64). Chronically elevated levels of IL-6 (> 2.0 ng/L) of middle-aged persons reduce the chance of successful ageing and increase the risk of future cardiovascular events or non-cardiovascular death later in life (1). Up to now it is unclear whether the main source of these pro-inflammatory factors is the chronically activated immune system (inflamm-ageing) or the senescence of cells with their senescence-associated secretory phenotype (SASP) leading to an enhanced secretion of pro-inflammatory mediators (15, 16, 18, 27).

Corresponding Author:

Ass.-Prof. DI Dr. Barbara Wessner, Centre for Sport Science and University Sports, University of Vienna, Auf der Schmelz 6, A-1150 Vienna, AUSTRIA
Tel.: +43-1-4277-28772, Fax: +43-1-4277-9287
E-Mail: barbara.wessner@univie.ac.at

Besides being associated with age-related diseases such as diabetes mellitus type 2, cardiac illnesses or neurological diseases, chronic subclinical inflammation may also contribute to impaired physical function in older adults. Several large scale studies have revealed that higher levels of circulating CRP or IL-6 are associated with lower physical performance such as handgrip strength or gait speed (4, 7, 12, 51). While alterations in TNF- α , IL-6 or CRP in response to ageing, physical inactivity as well as acute and chronic exercise have been studied extensively (14, 38, 42), much less is known about the involvement of transforming growth factor- β (TGF- β) in this context.

TGF- β is known as a potent regulatory cytokine with diverse effects on haemopoietic cells. It has pivotal function in the immune system through keeping tolerance via the regulation of lymphocyte proliferation, differentiation and survival (28). Furthermore, TGF- β controls the initiation and resolution of inflammatory responses through the regulation of chemotaxis, activation and survival of lymphocytes, natural killer cells, dendritic cells, macrophages, mast cells and granulocytes (31). It has been suggested that the age-associated dysregulation of the immune system might be caused by a decreased expression and functioning of receptors or signalling rafts and/or defects in signalling pathways finally leading to altered function of immune cells and immunosenescence (25). TGF- β mediates its biological function via binding to type II and forming a complex with type I transmembrane serine/threonine kinase receptors (TGF- β R1, TGF- β R2). Ligand binding assembles a complex consisting of two type I receptor components and two type II components, whereby the type I components are phosphorylated further activating intracellular Smad proteins whose nuclear localization is required for the transcriptional regulation of target genes (35).

Interestingly, TGF- β signalling is intimately involved in biogenesis and control of microRNAs (miRNAs) as Smad proteins play a regulatory role in the processing of miRNAs in the nucleus (2, 21). MiRNAs are small non-coding RNAs that affect gene expression either by inhibiting translation of their target proteins or by degrading the respective mRNA (67). They have been shown to regulate many biological functions including ageing, immune function and the response to exercise (33, 36, 53, 63). It also has been suggested that senescent donor cells contribute to SASP by secreting not only soluble proteins but also microvesicles containing miRNAs which are taken up by recipient cells and may cause or contribute to age related pathologies like osteoporosis, atherosclerosis, Alzheimer's disease or diabetes mellitus, type 2 (62). Several miRNAs such as miRNA-155, -146a and -21 are involved in the regulation of inflammation by controlling Toll-like receptor or nuclear factor- κ B (NF- κ B) signalling (45, 48). MiRNA-21, despite being well established as an 'onco-miR' due to its aberrant expression in numerous cancers, seems to be of particular importance as it links inflamm-ageing to cellular senescence (44). Intracellular as well as circulating miRNA-21 levels are higher in elderly in comparison to young subjects or centenarians. This is associated with a lower TGF- β R2 expression in leukocytes (46), whereby TGF- β R2 is a validated target of miRNA-21 in adipocytes (24) and colon cancer cells (66). Mice lacking the TGF- β R2 develop an

autoimmune biliary ductular disease similar to human primary biliary cirrhosis. Using this experimental model it has been shown that the lack of TGF- β signalling leads to a down-regulation of several miRNAs in T cells but interestingly to an up-regulation of miR-21 concomitant with an increased production of TNF- α and interferon- γ (2). It is hypothesized that the up-regulation of miRNA-21 is caused by a dysregulated gene expression normally controlled by TGF- β . This would result in the activation of inflammatory pathways such as NF- κ B which directly or indirectly (via induction of miRNA-21) leads to an up-regulation of inflammation. Another possibility is that a global down-regulation of miRNA expression induces multiple genes causing inflammation; this in turn could lead to elevated levels of miR-21 with a feedback up-regulation of inflammation (2).

However, TGF- β signalling seems to play an important role in relation to several diseases associated with chronic-low grade inflammation. We hypothesized that both, age and physical fitness could affect TGF- β and its receptors in peripheral blood mononuclear cells of healthy women. Therefore, the aim of the current study was to investigate (i) whether circulating TGF- β , as well as intracellular TGF- β , TGF- β receptors and miRNA-21 would differ between young and elderly women and (ii) whether these markers would be associated with parameters of physical performance.

METHODS

Subjects

Ninety elderly women (aged 65-92 years) who were recruited in 5 different senior residencies in the area of Vienna (Cura-torship of Viennese Retirement Homes (KWP)) participated in the study. For the current study we used baseline characteristics of study participants intended to take part in a prospective training study. In addition 16 young women (aged 18-28 years) responded to flyers at the University of Vienna. Young as well as elderly women were sedentary (less than 1h of sports activities per week) and free of severe diseases that would contra-indicate medical training therapy or measurement of physical performance, serious cardiovascular diseases, diabetic retinopathy and regular use of cortisone-containing drugs. Written informed consent was obtained from all participants before entry into the study in accordance with the Declaration of Helsinki and after approval by the ethics committee of the City of Vienna (EK-11-151-0811).

Anthropometric Measurements

Using a commercial stadiometer (Seca, Hamburg, Germany), standing height was measured without shoes to the nearest 0.5 cm. Shoulders kept in a relaxed position and arms allowed to hang freely. Body mass was evaluated with a digital scale (BWB 700, Tanita, Amsterdam, Netherlands) to the nearest 0.1 kg with subjects lightly dressed and barefoot. Body mass index (BMI) was calculated by dividing body mass in kilograms by height in meters squared. For determining body composition (muscle and fat mass) we used bioelectric impedance analyses, due to successful validation against data obtained by magnetic resonance imaging (50). Bioelectric Impedance Analyses (BIA) were performed in the morning

after an overnight fast using a BIA Analyzer 2000-S (Data-Input GmbH, Darmstadt, Germany). Participants were asked not to perform any exercise or strenuous physical activity the day before the tests.

Determination of physical performance

To evaluate each participant's aerobic endurance a *6-minutes walking test* was conducted. Therefore, participants walked for 6 minutes as fast as possible on a 30 metre shuttle track. They were allowed to reduce their speed or to rest if the selected speed was too high to be sustained. The completed distance within 6 minutes was recorded (55).

To measure *handgrip strength* participants performed two trials of an isometric handgrip strength test (kg) using a dynamometer in a sitting position with an angle of 45° in the elbow and the lower arm on the armrest. The participant was instructed to squeeze the handle as hard as possible for 4-5 seconds and the maximum isometric contraction was recorded (SAEHAN Corporation, Masan, Korea). The two trials were separated by one minute of passive recovery (37). Out of the two trials on each arm, the best result regardless of side was used for further calculation.

The *isokinetic peak torque of knee extension and flexion* consisted of concentric isokinetic torque measurements (Lido Loredan Biomedical, Inc., Davis, USA; Range of Motion 30°-80°, speed 60°/s or 120°/s). The left leg was tested in all participants except for 3 older women with acute injuries of the left leg making it necessary to test the right leg. The best result of two trials separated by a rest period of two minutes between the attempts was documented. Absolute values were divided by body mass to obtain relative values.

Blood sampling and analyses

Routine blood analyses

Between 06:30 and 08:00 in the morning venous blood samples were taken after an overnight fast. Venous blood was collected in Z Serum Clot Activator collection tubes (Vacuette®, Greiner Bio-One GmbH, Kremsmünster, Austria) for cytokine analyses and in EDTA tubes for the determination of leukocyte subpopulation numbers. For the isolation of peripheral blood mononuclear cells (PBMCs) from whole-blood, BD Vacutainer® CPT™ Tubes containing ~130 IU Na-Heparin and 2 ml FicoII™ (Becton, Dickinson and Company, Schwechat, Austria) were used.

After at least 30 min and at most 60 min after blood collection, the serum tubes were centrifuged (10 min, 3,000 x g). An aliquot of 1 ml was used for immediate determination of glucose, insulin and hs-CRP. The remaining serum was stored in aliquots at -80°C until further analysis. Glucose was analyzed by hexokinase method and insulin was estimated using a solid-phase, enzyme-labeled chemiluminescent immunometric assay (IMMULITE 2000, Siemens Healthcare Diagnostics Inc., Llanberis, UK). Cholesterol, HDL cholesterol, LDL cholesterol, triglyceride and hs-CRP were routinely quantified on a Cobas 8000 (Roche Diagnostics, Vienna, Austria). Leukocytes, lymphocytes, monocytes and granulocytes were quantified by flow cytometry on a Sysmex XE-2100™ Automated Hematology System (Sysmex Austria GmbH, Vienna, Austria).

Serum levels of TGF- β

TGF- β was determined using a commercially available DuoSet development kit for performing enzyme-linked immunosorbent assays (DY240, R&D Systems; Abingdon, UK) consisting of a capture antibody (2 μ g/ml of mouse anti-TGF- β 1), a detection antibody (300 ng/ml of biotinylated chicken anti-human TGF- β 1), and recombinant human TGF- β 1 to prepare a standard curve (31-2,000pg/ml). Twenty μ l of each serum sample were activated by adding 10 μ l of 1N HCl, incubated at room temperature for 10 min and neutralized with 10 μ l of 1.2N NaOH/0.5 M Hepes. The activated sample was diluted 20-fold with reagent diluent (0.05% Tween® 20 in PBS) and used in the assay following the instructions of the manufacturer. Spectrophotometric measurements were performed on a Victor³ 1420 Multilabel Counter (Perkin Elmer, MA, US).

Isolation of total RNA from PBMCs

PBMCs were separated from red blood cells and neutrophils by centrifugation of BD Vacutainer® CPT Tubes at 1,650 x g for 20 min at room temperature. After removing 2 ml of the plasma supernatant, the cells comprising PBMCs were resuspended in the remaining plasma and transferred to another tube. PBMCs were washed twice with PBS without Ca and Mg according to the protocol provided by the manufacturer. Finally, the pellet was carefully resuspended in 700 μ l of QIAzol Lysis Reagent (Qiagen, Hilden, Germany) and stored on -80° until analysis.

Total RNA including small RNAs was isolated after thawing and incubating the samples for 5 min at room temperature using the miRNeasy Mini Kit (Qiagen, Hilden, Germany) and following the instructions of the manufacturer. In order to prepare a miRNA-enriched fraction separated from the larger RNAs (>200nt) the RNeasy Min Elute Cleanup Kit (Qiagen, Hilden, Germany) was used. Reverse transcription was for the miRNA-enriched fraction was performed using the miScript II RT Kit (Qiagen, Hilden, Germany) while larger RNAs were reverse transcribed using the QuantiTect Reverse Transcription Kit (Qiagen, Hilden, Germany).

Quantitative real-time RT-PCR

TGF- β , TGF- β RI and TGF- β RII mRNA were determined using the respective primer assays (Hs_TGFB1_1 (QT00000728), Hs_TGFB1_1 (QT00083412), Hs_TGFB2_1 (QT00014350), Qiagen, Hilden, Germany) in conjunction with the QuantiTect SYBR Green PCR kit (Qiagen, Hilden, Germany). A standard curve was prepared by pooling equal amounts of cDNA from PBMCs of 7 young and 23 old subjects which were randomly selected from the study population. In addition, GAPDH (Hs_GAPDH_2 (QT01192646)) served as endogenous control and was used to normalize the data. Quantification was performed on an Applied Biosystems® 7500 Real-Time PCR System.

MiRNA-21 expression levels were detected using a miScript Primer Assay specific for miRNA-21 (hs_miR-21_2 (MS00009079), Qiagen, Hilden, Germany). A standard curve was prepared by using a commercially available totalRNA of peripheral blood leukocyte cells of a 24 year old female donor (Total RNA (R1234148-10), BioChain, Newark, USA). Quan-

tification was performed on an Applied Biosystems® 7500 Real-Time PCR System.

Statistical analyses

The data acquisition and data processing took place using commercial software (IBM SPSS for Windows, Version 20). Shapiro-Wilk test was used to determine if data sets were normally distributed. As most of the variables did not meet the criteria for normal distribution, Mann-Whitney U test was used in order to compare young and elderly women. Associations between variables were analysed using Spearman's ρ correlation coefficient. Data are shown as median (minimum-maximum), statistical significance was set at $p < 0.05$.

RESULTS

Subject characteristics

Old females with a median age of 84 years had a higher body mass (+23%, $p < 0.001$) as well as BMI (+36%, $p < 0.001$) compared to young females (median age: 25 years). Bioelectric impedance analyses revealed a higher percentage of body fat mass (+37%, $p < 0.001$) and a lower muscle mass (-12%, $p < 0.001$). The age-related changes in body composition were accompanied by substantial worsening of health-related parameters such as circulating glucose levels (+10%, $p < 0.001$), cholesterol (+22%, $p = 0.002$), LDL cholesterol (+49%, $p < 0.001$), HDL cholesterol (-16%, $p = 0.006$), and triglyceride (+41%, $p = 0.006$) (Table 1).

Physical fitness

Aerobic fitness was determined by 6-minutes walking test (6MWT). Old females reached a significant lower distance within a time frame of 6 minutes (-61%, $p < 0.001$) than younger females (Figure 1A). Strength was determined by handgrip dynamometer as well as isokinetic peak torque measurements. Isometric handgrip strength was significantly lower in old women (-43%, $p < 0.001$, Figure 1B). Similarly, relative peak torque knee extension (PTE) as well as relative peak torque knee flexion (PTF) differed significantly between groups at both tested velocities (Fig. 1C-F) reflecting strength loss of M. quadriceps (-48% at 60°/s and -51% at 120°/s, $p < 0.001$) and hamstrings (-53% at 60°/s as well as at 120°/s, $p < 0.001$).

Inflammatory parameters

Differences in inflammatory parameters between young and old females are summarized in Table 2. The number of leukocytes in whole blood did not differ between young and old women. However, the subpopulation analysis revealed a significant higher percentage of monocytes in elderly (+24%, $p = 0.007$), with no changes in lymphocytes and granulocytes. As expected, hs-CRP was significantly higher in old women in comparison to young females (+156%, $p < 0.001$). Serum levels of TGF- β did not differ between young and elderly women ($p = 0.290$). Similarly, TGF- β mRNA levels from PBMCs did not vary between groups ($p = 0.290$). Interestingly, both TGF- β RI (-49%, $p = 0.020$) and TGF- β RII mRNA levels (-46%, $p = 0.020$) were lower with respect to old age.

Table 1: Subject Characteristics

Parameter	Young (n=16)	Old (n=90)	<i>p-value</i>
Age [years]	24.9 (21.7-28.4)	83.8 (65.0-92.2)	<0.001
Body mass [kg]	58.1 (51.0-65.2)	71.7 (46.2-112.4)	<0.001
Height [m]	1.65 (1.57-1.71)	1.57 (1.40-1.72)	<0.001
BMI [kg/m ²]	21.7 (18.9-23.6)	29.6 (18.1-50.0)	<0.001
Body fat mass [%]	26.3 (21.2-33.2)	36.1 (14.0-50.4)	<0.001
Muscle mass [kg]	22.2 (20.6-23.1)	19.5 (12.9-26.4)	0.019
Glucose [mg/dl]	87 (67-106)	96 (79-196)	<0.001
Insulin [μ IU/ml]	6.40 (2.50-13.50)	8.05 (1.32-41.57)	0.068
Cholesterol [mg/dl]	171 (136-264)	209 (144-336)	0.002
HDL-Cholesterol [mg/dl]	74 (54-99)	62 (33-120)	0.006
LDL-Cholesterol [mg/dl]	81 (47-146)	121 (43-238)	<0.001
Triglyceride [mg/dl]	79 (41-191)	111 (43-275)	0.006

Data are expressed as medians (min-max); Differences were detected using Mann-Whitney U test; BMI (body mass index); HDL (high density lipoprotein); LDL (low density lipoprotein)




Figure 1. Age-related differences in physical fitness: (A) 6-minutes Walking Test, (B) Handgrip strength, (C) Relative Peak Torque Knee Extension (PTE) 60°/s, (D) Relative Peak Torque Knee Flexion (PTF) 60°/s, (E) Relative Peak Torque Knee Extension (PTE) 120°/s, (F) Relative Peak Torque Knee Flexion (PTF) 120°/s. # denotes a significant differences between young and old women ($p < 0.001$).

To investigate whether the change in TGF- β receptor mRNA expression could be influenced by miRNA-21, its level was measured in PBMCs. While we did not detect any differences in intracellular miRNA-21 levels between young and old women ($p = 0.190$), miRNA-21 in PBMCs of old women correlated negatively with TGF- β RI expression ($\rho = -0.315$; $p = 0.004$) but not with TGF- β RII or TGF- β .

Correlations between fitness and inflammation in elderly women

Next we were interested whether higher fitness levels within the cohort of older women would be associated with lower pro-inflammatory states (Table 3). Indeed, hs-CRP levels and relative peak torque measurements revealed negative correlation; however, significance was reached only for relative peak torque of knee extension at 120°/s ($\rho = -0.276$; $p = 0.013$). Surprisingly, hs-CRP was not associated with performance in 6MWT and even positively correlated to handgrip strength

($\rho = 0.237$; $p = 0.035$). Serum TGF- β , its expression level in PBMCs, and the expression of its receptors TGF- β RI and TGF- β RII were not related to any performance measure.

Body composition influences inflammatory and fitness-related parameters in elderly

As the subgroup of older women showed a substantial variation in body composition, its influence on inflammatory, fitness- and health-related variables was investigated. It has to be mentioned that age in this subgroup was even negatively associated with muscle mass ($\rho = -0.546$; $p < 0.001$) but surprisingly also with BMI ($\rho = -0.219$; $p = 0.039$) and body fat mass ($\rho = -0.316$; $p = 0.003$).

Hs-CRP was positively correlated to BMI ($\rho = 0.326$; $p = 0.002$), body fat ($\rho = 0.331$; $p = 0.002$) and muscle mass ($\rho = 0.291$; $p = 0.007$), but not age. Furthermore, total leukocyte counts were positively associated with BMI ($\rho = 0.342$; $p = 0.001$) and body fat ($\rho = 0.212$; $p = 0.050$). In contrast, TGF-

Table 2: Age-related changes in immune parameters

Parameter	Young (n=16)	Old (n=90)	p-value
<u>Circulating</u>			
Leukocytes [$10^9/l$]	6.0 (4.6-8.0)	6.5 (3.4-13.3)	0.208
Lymphocytes [%]	33.8 (23.5-46.6)	31.6 (15.3-48.1)	0.375
Monocytes [%]	6.7 (3.7-11.5)	8.3 (1.0-14.1)	0.007
Granulocytes [%]	59.0 (43.8-67.2)	56.3 (38.4-75.0)	0.584
hs-CRP [mg/l]	0.9 (0.1-4.3)	2.3 (0.3-56.7)	0.001
TGF- β [pg/ml]	33,321 (23,793-42,543)	34,851 (16,667-73,681)	0.290
<u>PBMCs (intracellular)</u>			
TGF- β / GAPDH [-]	0.65 (0.30-1.59)	0.53 (0.06-3.36)	0.387
TGF- β RI / GAPDH [-]	4.07 (1.38-14.60)	2.08 (0.14-28.81)	0.022
TGF- β RII / GAPDH [-]	3.16 (1.14-10.25)	1.71 (0.51-14.86)	0.022
hsa-miRNA-21 [copies /pg RNA]	1,821 (380-3,824)	2,452 (57-5,481)	0.190

Data are expressed as medians (min-max); Differences were detected using Mann-Whitney U test; hs-CRP (high sensitive C-reactive protein), PBMC (peripheral blood mononuclear cells), TGF- β (transforming growth factor- β), TGF- β RI (transforming growth factor- β receptor type I), TGF- β RII (transforming growth factor- β receptor type II), GAPDH (glyceraldehyde-3-phosphate dehydrogenase); hsa-miRNA-21 (human microRNA-21)

Table 3: Correlation between fitness and immune parameters in old women

	6MWT	Handgrip	PTE _{rel} 60°/s	PTF _{rel} 60°/s	PTE _{rel} 120°/s	PTF _{rel} 120°/s
Leukocytes	-0.100	-0.102	-0.017	0.062	-0.020	-0.039
hs-CRP	0.049	0.237*	-0.206	-0.127	-0.276*	-0.175
TGF-β (circulating)	-0.086	-0.082	-0.060	-0.019	-0.046	-0.120
TGF-β / GAPDH (intracellular)	-0.117	-0.043	0.033	-0.101	0.138	0.033
TGF-βRI / GAPDH (intracellular)	-0.070	-0.092	-0.124	-0.180	-0.013	-0.089
TGF-βRII / GAPDH (intracellular)	-0.175	-0.045	-0.002	-0.116	0.092	0.038
hsa-miRNA-21 (intracellular)	-0.011	0.243*	0.101	0.034	0.075	0.015

Data indicate Spearman-Rho correlation coefficients. * $p < 0.05$, $n \geq 80$; 6MWT (6 Minutes Walking Test), PTE_{rel} (relative peak torque of knee extension), PTF_{rel} (relative peak torque of knee flexion), hs-CRP (high sensitive C-reactive protein), TGF- β (transforming growth factor- β), TGF- β RI (transforming growth factor- β receptor type I), TGF- β RII (transforming growth factor- β receptor type II), GAPDH (glyceraldehyde-3-phosphate dehydrogenase), hsa-miRNA-21 (human microRNA-21)

β RII mRNA was negatively associated with body fat ($\rho = -0.263$, $p = 0.018$) but not with BMI ($\rho = -0.185$; $p = 0.091$). None of the other inflammatory variables correlated with body composition.

With respect to fitness parameters BMI and body fat correlated negatively with relative peak torque of knee extension at 120°/s (BMI: $\rho = -0.282$; $p = 0.011$, body fat: $\rho = -0.237$; $p = 0.038$) and partly with relative peak torque of knee flexion

at 120°/s (BMI: $\rho=-0.292$; $p=0.009$, body fat: $\rho=-0.209$; $p=0.068$). BMI and body mass were not associated with handgrip strength or 6MWT. Muscle mass correlated positively with handgrip strength ($\rho=0.662$; $p<0.001$), but not with other performance parameters.

DISCUSSION

The aim of the current study was to investigate the expression of TGF- β , its receptors and its potential modulator miRNA-21 in PBMCs in the context of age and fitness status. Young and old females differed substantially in fitness as measured by 6MWT, handgrip strength, isokinetic peak torque of knee extensors and flexors as well as serum hs-CRP levels. While serum levels of TGF- β as well as TGF- β and miRNA-21 expression levels in PBMCs did not differ between young and old females, TGF- β R1 and TGF- β R2 mRNA were significantly lower in the elderly. However, within the cohort of elderly women neither TGF- β nor its receptors were associated with performance characteristics.

Initially, TGF- β was purified from human platelets (3). In mammals three different isoforms (TGF- β 1, - β 2, - β 3) have been described, whereby TGF- β 1 is the predominant form in immune cells. TGF- β is synthesized and secreted by most cell types as an inactive precursor complex, termed latent TGF- β , where TGF- β is bound non-covalently to the latency associated peptide (LAP). To be activated TGF- β has to be cleaved from the LAP using one of physiological mechanisms such as proteolytic cleavage by plasmin, cathepsin, and other enzymes, oxidation by free radicals or the interaction with thrombospondin (41). Circulating TGF- β has been detected in a variety of studies with plasma values ranging from below 0.1 ng/ml up to more than 25 ng/ml. Of course the study population differed between these studies ranging from healthy individuals of mixed gender and age-groups to different patient groups. However, methodological issues concerning plasma processing and assay system seem to be the most critical factors in determining TGF- β levels (20). In this respect it has been shown that platelet degranulation, haemolysis of erythrocytes and contamination with leukocytes can lead to an over-estimation of TGF- β protein levels in plasma, however different activation protocols to dissociate TGF- β from its complexes are in use. For this purpose, we and many other groups used acidification of serum samples prior to assessing TGF- β by ELISA (20). Furthermore, both serum from young and old females were treated in the same way, therefore minimizing the risk of bias within the study.

Regarding lifestyle-related diseases TGF- β seems to play conflicting roles as shown for cancer, where it can either act as tumour suppressor by inhibiting cell proliferation and inflammation in the early stage of cancer development or as tumour promoter by inducing metastasis or angiogenesis in later stages of the disease (52). Similarly, higher levels of TGF- β are measured in hypertensive humans (57), but increased serum levels of TGF- β may also protect patients with coronary artery disease against cardiovascular events and coronary interventions (59). With respect to age decreased levels have been reported for adults (21-67 years)

in comparison to children (1-14 years) (43). Another study has revealed that TGF- β levels are higher in males than in females, but decrease with age and increase with obesity in both genders (32). However, in centenarians, serum TGF- β concentration seems to be higher than in younger adults, suggesting that high concentrations of TGF- β might be beneficial during extreme old age (6). These data are in conflict with our results as we did not detect any differences in TGF- β between young and old women. However, the broad range of age as well as other lifestyle related factors such as obesity or diabetes which were characteristic for our study could have influenced the results.

Conflicting data have also been reported with regard to TGF- β and acute exercise, whereby intensity and type of exercise seem to play an important role for data interpretation. While a graded cycling exercise to exhaustion of about 18 min duration (10) and 1 h of treadmill running at about 70-80% of VO_{2max} (22) are able to increase the concentration of circulating TGF- β , 1 h of cycling exercise at $\sim 70\%$ of VO_{2max} does not alter circulating TGF- β (17). However, salivary TGF- β is increased as late as 24 h after a moderate exercise bout (49). Long-term training for 6 weeks in healthy students resulted in a biphasic response of TGF- β with increased levels after 2 weeks of training and lower levels at the end of the training period (23). This is in contrast to another study in diabetic patients showing that 8 weeks of strength and aerobic training results in increased TGF- β levels which are accompanied by lower hs-CRP levels (61). Furthermore, 6 months of exercise (2.5 h per week) were able to increase TGF- β production of unstimulated as well as phytohaemagglutinin-stimulated peripheral blood mononuclear cells of persons at risk of developing ischemic heart disease (54). Taken together, it seems that chronic exercise lowers TGF- β in young and healthy persons but it might up-regulate TGF- β in patients suffering from lifestyle-related diseases.

In addition to circulating levels of TGF- β we were especially interested in expression of TGF- β and its receptors in PBMCs of young and old women. While intracellular TGF- β mRNA was not different between these two groups, lowered TGF- β R1 and TGF- β R2 have been detected in elderly. These results are partly in accordance with a previous study in young (20-30 years), old (75-85 years) and very old (>98 years) subjects, where leukocyte TGF- β R2 mRNA were lowest in the 75-85 year old group in comparison to both, the young and the centenarians. In contrast to our study the decrease in TGF- β R2 expression is accompanied by an increase in miRNA-21 levels. However, neither intracellular nor circulating TGF- β nor physical performance has been assessed in this study (46). The importance of signalling via TGF- β R2 has been shown in an animal model where the induction of a TGF- β R2 gene disruption results in a lethal inflammatory disease (29). On the one hand TGF- β R2 is important in T-cell mediated immunity (30) but on the other hand macrophages lacking TGF- β R2 have defects in expression of a set of genes that form the hallmark of the M2 polarizing program in macrophages which is important to induce the anti-inflammatory effects of M2 macrophages such as phagocytosis of apoptotic cells, resolution of inflammation and tissue repair (19, 34, 40).

Data support the hypothesis that exercise can reduce low grade inflammation in elderly (61) and provide long-term benefits with regard to cardiovascular, cognitive, psychosocial and other aspects in elderly (11). Hs-CRP has been shown to be consistently higher in elderly (26, 60), a fact that was confirmed in the current study. Moreover, hs-CRP correlated positively with BMI and body fat but negatively with relative peak torque measurements. This partly confirms several studies which revealed associations between a higher inflammatory state and lower physical performance (47, 56). Besides originating in the liver, the acute phase protein hs-CRP is produced and released from adipose tissue thereby linking obesity to a chronic inflammatory state (13, 65).

Although we detected a negative correlation between hs-CRP and isokinetic knee extension strength which is in line with many studies linking chronic inflammation to low physical performance in elderly (7, 9, 12, 58), this picture was not consistent for other performance parameters such as aerobic fitness or strength. The reason for this finding could be a complex interaction between several factors which has been suggested by Morrisette-Thomas et al. who applied principal component analysis in order to understand why inflamm-aging does not simply reflect increases in pro-inflammatory markers (39). One especially interesting aspect in our study was that general fitness status of elderly women was not related to TGF- β , TGF- β RI or TGF- β RII expression in PBMCs. Furthermore, higher hs-CRP and miRNA-21 levels in older women were even associated with a higher handgrip strength. According to our hypothesis we would have expected higher levels of the inflammatory miRNA-21 in subjects with low physical performance as suggested by Bye et al. who showed that miRNA-21 was increased in male participants with low aerobic capacity as assessed by VO₂max (5). However, these studies are sparsely comparable as miRNA-21 was detected in serum, participants were male and younger and there might be a difference between performance indicators for strength or endurance. Furthermore, it would have been interesting to measure general physical activity levels by an objective method such as accelerometry as both current physical activity practice and performance are associated with inflammatory biomarkers (12).

In summary, it has been shown that TGF- β is involved in a variety of physiological as well as pathological process exerting positive but in some cases also negative effects on health. We have demonstrated that in older women TGF- β signalling in PBMCs might be impaired as reflected by a lower gene expression of its receptors TGF- β RI and TGF- β RII independent of physical fitness. However, further studies are needed to test whether a reduced expression of the TGF- β receptors indeed would reduce TGF- β signalling in PBMCs in order to get mechanistic insight as well as to reveal its functional consequences, more precisely.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the Curatorship of Viennese retirement home (KWP) and residents for taking part in the study. Financial support was provided by the University of Vienna by funding the Research Platform Active Ageing.

REFERENCES

1. Akbaraly TN, Hamer M, Ferrie JE, Lowe G, Batty GD, Hagger-Johnson G, Singh-Manoux A, Shipley MJ and Kivimaki M. Chronic inflammation as a determinant of future aging phenotypes. *CMAJ* 185(16):E763-770, 2013.
2. Ando Y, Yang GX, Kenny TP, Kawata K, Zhang W, Huang W, Leung PS, Lian ZX, Okazaki K, Ansari AA, He XS, Invernizzi P, Ridgway WM, Lu Q and Gershwin ME. Overexpression of microRNA-21 is associated with elevated pro-inflammatory cytokines in dominant-negative TGF-beta receptor type II mouse. *J Autoimmun* 41:111-119, 2013.
3. Assoian RK, Komoriya A, Meyers CA, Miller DM and Sporn MB. Transforming growth factor-beta in human platelets. Identification of a major storage site, purification, and characterization. *J Biol Chem* 258(11):7155-7160, 1983.
4. Brinkley TE, Leng X, Miller ME, Kitzman DW, Pahor M, Berry MJ, Marsh AP, Kritchevsky SB and Nicklas BJ. Chronic inflammation is associated with low physical function in older adults across multiple comorbidities. *J Gerontol A Biol Sci Med Sci* 64(4):455-461, 2009.
5. Bye A, Rosjo H, Aspenes ST, Condorelli G, Omland T and Wisloff U. Circulating microRNAs and aerobic fitness--the HUNT-Study. *PLoS One* 8(2):e57496, 2013.
6. Carrieri G, Marzi E, Olivieri F, Marchegiani F, Cavallone L, Cardelli M, Giovagnetti S, Stecconi R, Molendini C, Trapassi C, De Benedictis G, Klekas D and Franceschi C. The G/C915 polymorphism of transforming growth factor beta1 is associated with human longevity: a study in Italian centenarians. *Aging Cell* 3(6):443-448, 2004.
7. Cesari M, Penninx BW, Pahor M, Lauretani F, Corsi AM, Rhys Williams G, Guralnik JM and Ferrucci L. Inflammatory markers and physical performance in older persons: the InCHIANTI study. *J Gerontol A Biol Sci Med Sci* 59(3):242-248, 2004.
8. Chung HY, Sung B, Jung KJ, Zou Y and Yu BP. The molecular inflammatory process in aging. *Antioxid Redox Signal* 8(3-4):572-581, 2006.
9. Colbert LH, Visser M, Simonsick EM, Tracy RP, Newman AB, Kritchevsky SB, Pahor M, Taaffe DR, Brach J, Rubin S and Harris TB. Physical activity, exercise, and inflammatory markers in older adults: findings from the Health, Aging and Body Composition Study. *J Am Geriatr Soc* 52(7):1098-1104, 2004.
10. Czarkowska-Paczek B, Bartlomiejczyk I and Przybylski J. The serum levels of growth factors: PDGF, TGF-beta and VEGF are increased after strenuous physical exercise. *J Physiol Pharmacol* 57(2):189-197, 2006.
11. de Gonzalo-Calvo D, de Luxan-Delgado B, Martinez-Cambor P, Rodriguez-Gonzalez S, Garcia-Macia M, Suarez FM, Solano JJ, Rodriguez-Colunga MJ and Coto-Montes A. Chronic inflammation as predictor of 1-year hospitalization and mortality in elderly population. *Eur J Clin Invest* 42(10):1037-1046, 2012.

12. Elosua R, Bartali B, Ordovas JM, Corsi AM, Lauretani F, Ferrucci L and In CI. Association between physical activity, physical performance, and inflammatory biomarkers in an elderly population: the InCHIANTI study. *J Gerontol A Biol Sci Med Sci* 60(6):760-767, 2005.
13. Farhangi MA, Keshavarz SA, Eshraghian M, Ostadrahimi A and Saboor-Yaraghi AA. White blood cell count in women: relation to inflammatory biomarkers, haematological profiles, visceral adiposity, and other cardiovascular risk factors. *J Health Popul Nutr* 31(1):58-64, 2013.
14. Fischer CP, Berntsen A, Perstrup LB, Eskildsen P and Pedersen BK. Plasma levels of interleukin-6 and C-reactive protein are associated with physical inactivity independent of obesity. *Scand J Med Sci Sports* 17(5):580-587, 2007.
15. Franceschi C, Capri M, Monti D, Giunta S, Olivieri F, Sevini F, Panourgia MP, Invidia L, Celani L, Scurti M, Cevenini E, Castellani GC and Salvioli S. Inflammaging and anti-inflammaging: a systemic perspective on aging and longevity emerged from studies in humans. *Mech Ageing Dev* 128(1):92-105, 2007.
16. Freund A, Orjalo AV, Desprez PY and Campisi J. Inflammatory networks during cellular senescence: causes and consequences. *Trends Mol Med* 16(5):238-246, 2010.
17. Garcia JJ, Bote E, Hinchado MD and Ortega E. A single session of intense exercise improves the inflammatory response in healthy sedentary women. *J Physiol Biochem* 67(1):87-94, 2011.
18. Giunta S. Exploring the complex relations between inflammation and aging (inflamm-aging): anti-inflamm-aging remodeling of inflamm-aging, from robustness to frailty. *Inflamm Res* 57(12):558-563, 2008.
19. Gong D, Shi W, Yi SJ, Chen H, Groffen J and Heisterkamp N. TGFbeta signaling plays a critical role in promoting alternative macrophage activation. *BMC Immunol* 13:31, 2012.
20. Grainger DJ, Mosedale DE and Metcalfe JC. TGF-beta in blood: a complex problem. *Cytokine Growth Factor Rev* 11(1-2):133-145, 2000.
21. Hata A and Davis BN. Control of microRNA biogenesis by TGFbeta signaling pathway-A novel role of Smads in the nucleus. *Cytokine Growth Factor Rev* 20(5-6):517-521, 2009.
22. Heinemeier K, Langberg H and Kjaer M. Exercise-induced changes in circulating levels of transforming growth factor-beta-1 in humans: methodological considerations. *Eur J Appl Physiol* 90(1-2):171-177, 2003.
23. Hering S, Jost C, Schulz H, Hellmich B, Schatz H and Pfeiffer H. Circulating transforming growth factor beta1 (TGFbeta1) is elevated by extensive exercise. *Eur J Appl Physiol* 86(5):406-410, 2002.
24. Kim YJ, Hwang SJ, Bae YC and Jung JS. MiR-21 regulates adipogenic differentiation through the modulation of TGF-beta signaling in mesenchymal stem cells derived from human adipose tissue. *Stem Cells* 27(12):3093-3102, 2009.
25. Kohut ML and Senchina DS. Reversing age-associated immunosenescence via exercise. *Exerc Immunol Rev* 10:6-41, 2004.
26. Kritchevsky SB, Cesari M and Pahor M. Inflammatory markers and cardiovascular health in older adults. *Cardiovasc Res* 66(2):265-275, 2005.
27. Kuilman T and Peeper DS. Senescence-messaging secretome: SMS-ing cellular stress. *Nat Rev Cancer* 9(2):81-94, 2009.
28. Le Blanc I, Laurent M, Bokobza B, Michot F and Teniere P. [Carney's triad: a new case associated with adrenal cortex adenoma]. *Gastroenterol Clin Biol* 14(4):399-401, 1990.
29. Leveen P, Larsson J, Ehinger M, Cilio CM, Sundler M, Sjöstrand LJ, Holmdahl R and Karlsson S. Induced disruption of the transforming growth factor beta type II receptor gene in mice causes a lethal inflammatory disorder that is transplantable. *Blood* 100(2):560-568, 2002.
30. Li MO, Sanjabi S and Flavell RA. Transforming growth factor-beta controls development, homeostasis, and tolerance of T cells by regulatory T cell-dependent and -independent mechanisms. *Immunity* 25(3):455-471, 2006.
31. Li MO, Wan YY, Sanjabi S, Robertson AK and Flavell RA. Transforming growth factor-beta regulation of immune responses. *Annu Rev Immunol* 24:99-146, 2006.
32. Lin Y, Nakachi K, Ito Y, Kikuchi S, Tamakoshi A, Yagyu K, Watanabe Y, Inaba Y, Kazuo T and Jacc Study G. Variations in serum transforming growth factor-beta1 levels with gender, age and lifestyle factors of healthy Japanese adults. *Dis Markers* 27(1):23-28, 2009.
33. Makarova JA, Maltseva DV, Galatenko VV, Abbasi A, Maximenko DG, Grigoriev AI, Tonevitsky AG and Northoff H. Exercise Immunology Meets MiRNAs. *Exerc Immunol Rev* 20:135-164, 2014.
34. Martinez FO, Helming L and Gordon S. Alternative activation of macrophages: an immunologic functional perspective. *Annu Rev Immunol* 27:451-483, 2009.
35. Massague J. TGFbeta signalling in context. *Nat Rev Mol Cell Biol* 13(10):616-630, 2012.
36. McGregor RA, Poppitt SD and Cameron-Smith D. Role of microRNAs in the age-related changes in skeletal muscle and diet or exercise interventions to promote healthy aging in humans. *Ageing Res Rev*, 2014.
37. Mijnders DM, Meijers JM, Halfens RJ, ter Borg S, Luiking YC, Verlaan S, Schoberer D, Cruz Jentoft AJ, van Loon LJ and Schols JM. Validity and reliability of tools to measure muscle mass, strength, and physical performance in community-dwelling older people: a systematic review. *J Am Med Assoc* 14(3):170-178, 2013.
38. Mikkelsen UR, Couppe C, Karlsen A, Grosset JF, Schjerling P, Mackey AL, Klausen HH, Magnusson SP and Kjaer M. Life-long endurance exercise in humans: Circulating levels of inflammatory markers and leg muscle size. *Mech Ageing Dev* 134(11-12):531-540, 2013.
39. Morrisette-Thomas V, Cohen AA, Fulop T, Riesco E, Legault V, Li Q, Milot E, Dusseault-Belanger F and Ferrucci L. Inflamm-aging does not simply reflect increases in pro-inflammatory markers. *Mech Ageing Dev* 139:49-57, 2014.
40. Mosser DM and Edwards JP. Exploring the full spectrum of macrophage activation. *Nat Rev Immunol* 8(12):958-969, 2008.
41. Murphy-Ullrich JE and Poczatek M. Activation of latent TGF-beta by thrombospondin-1: mechanisms and physiology. *Cytokine Growth Factor Rev* 11(1-2):59-69, 2000.
42. Neubauer O, König D and Wagner KH. Recovery after an Ironman triathlon: sustained inflammatory responses and muscular stress. *Eur J Appl Physiol* 104(3):417-426, 2008.
43. Okamoto Y, Gotoh Y, Uemura O, Tanaka S, Ando T and Nishida M. Age-dependent decrease in serum transforming growth factor (TGF)-beta 1 in healthy Japanese individuals; population study of serum TGF-beta 1 level in Japanese. *Dis Markers* 21(2):71-74, 2005.

44. Olivieri F, Rippo MR, Monsurro V, Salvioli S, Capri M, Procopio AD and Franceschi C. MicroRNAs linking inflammation, cellular senescence and cancer. *Ageing Res Rev* 12(4):1056-1068, 2013.
45. Olivieri F, Rippo MR, Procopio AD and Fazioli F. Circulating inflamma-miRs in aging and age-related diseases. *Front Genet* 4:121, 2013.
46. Olivieri F, Spazzafumo L, Santini G, Lazzarini R, Albertini MC, Rippo MR, Galeazzi R, Abbatecola AM, Marcheselli F, Monti D, Ostan R, Cevenini E, Antonicelli R, Franceschi C and Procopio AD. Age-related differences in the expression of circulating microRNAs: miR-21 as a new circulating marker of inflammaging. *Mech Ageing Dev* 133(11-12):675-685, 2012.
47. Omran Simin F, Narges Z, Sajad A, Parisa Y and Omrani Vahid F. Relationship between inflammatory markers of cardiovascular disease and VO₂peak in asymptomatic females. *J Sports Med Phys Fitness* 53(2):198-202, 2013.
48. Quinn SR and O'Neill LA. A trio of microRNAs that control Toll-like receptor signalling. *Int Immunol* 23(7):421-425, 2011.
49. Rosa L, Teixeira A, Lira F, Tufik S, Mello M and Santos R. Moderate acute exercise (70% VO₂ peak) induces TGF- β , alpha-amylase and IgA in saliva during recovery. *Oral Dis* 20(2):186-190, 2014.
50. Roubenoff R, Baumgartner RN, Harris TB, Dallal GE, Hannan MT, Economos CD, Stauber PM, Wilson PW and Kiel DP. Application of bioelectrical impedance analysis to elderly populations. *J Gerontol A Biol Sci Med Sci* 52(3):M129-136, 1997.
51. Sanders JL, Ding V, Arnold AM, Kaplan RC, Cappola AR, Kizer JR, Boudreau RM, Cushman M and Newman AB. Do changes in circulating biomarkers track with each other and with functional changes in older adults? *J Gerontol A Biol Sci Med Sci* 69(2):174-181, 2014.
52. Sheen YY, Kim MJ, Park SA, Park SY and Nam JS. Targeting the Transforming Growth Factor-beta Signaling in Cancer Therapy. *Biomol Ther (Seoul)* 21(5):323-331, 2013.
53. Smith-Vikos T and Slack FJ. MicroRNAs and their roles in aging. *J Cell Sci* 125(Pt 1):7-17, 2012.
54. Smith JK, Dykes R, Douglas JE, Krishnaswamy G and Berk S. Long-term exercise and atherogenic activity of blood mononuclear cells in persons at risk of developing ischemic heart disease. *JAMA* 281(18):1722-1727, 1999.
55. Steffen TM, Hacker TA and Mollinger L. Age- and gender-related test performance in community-dwelling elderly people: Six-Minute Walk Test, Berg Balance Scale, Timed Up & Go Test, and gait speeds. *Phys Ther* 82(2):128-137, 2002.
56. Stenholm S, Rantanen T, Heliovaara M and Koskinen S. The mediating role of C-reactive protein and handgrip strength between obesity and walking limitation. *J Am Geriatr Soc* 56(3):462-469, 2008.
57. Suthanthiran M, Li B, Song JO, Ding R, Sharma VK, Schwartz JE and August P. Transforming growth factor-beta 1 hyperexpression in African-American hypertensives: A novel mediator of hypertension and/or target organ damage. *Proc Natl Acad Sci U S A* 97(7):3479-3484, 2000.
58. Taaffe DR, Harris TB, Ferrucci L, Rowe J and Seeman TE. Cross-sectional and prospective relationships of interleukin-6 and C-reactive protein with physical performance in elderly persons: MacArthur studies of successful aging. *J Gerontol A Biol Sci Med Sci* 55(12):M709-715, 2000.
59. Tashiro H, Shimokawa H, Sadamatu K and Yamamoto K. Prognostic significance of plasma concentrations of transforming growth factor-beta in patients with coronary artery disease. *Coron Artery Dis* 13(3):139-143, 2002.
60. Topinkova E. Aging, disability and frailty. *Ann Nutr Metab* 52 Suppl 1:6-11, 2008.
61. Touvra AM, Volaklis KA, Spassis AT, Zois CE, Douda HD, Kotsa K and Tokmakidis SP. Combined strength and aerobic training increases transforming growth factor-beta1 in patients with type 2 diabetes. *Hormones (Athens)* 10(2):125-130, 2011.
62. Weilner S, Schraml E, Redl H, Grillari-Voglauer R and Grillari J. Secretion of microvesicular miRNAs in cellular and organismal aging. *Exp Gerontol* 48(7):626-633, 2013.
63. Wessner B, Gryadunov-Masutti L, Tschan H, Bachl N and Roth E. Is there a role for microRNAs in exercise immunology? A synopsis of current literature and future developments. *Exerc Immunol Rev* 16:22-39, 2010.
64. Woods JA, Wilund KR, Martin SA and Kistler BM. Exercise, inflammation and aging. *Ageing Dis* 3(1):130-140, 2012.
65. Wu T, Dorn JP, Donahue RP, Sempos CT and Trevisan M. Associations of serum C-reactive protein with fasting insulin, glucose, and glycosylated hemoglobin: the Third National Health and Nutrition Examination Survey, 1988-1994. *Am J Epidemiol* 155(1):65-71, 2002.
66. Yu Y, Kanwar SS, Patel BB, Oh PS, Nautiyal J, Sarkar FH and Majumdar AP. MicroRNA-21 induces stemness by downregulating transforming growth factor beta receptor 2 (TGFbetaR2) in colon cancer cells. *Carcinogenesis* 33(1):68-76, 2012.
67. Zhang R and Su B. Small but influential: the role of microRNAs on gene regulatory network and 3'UTR evolution. *J Genet Genomics* 36(1):1-6, 2009.